

White Paper on Non-O157:H7 Shiga-toxin Producing *E. coli* from Meat and Non-Meat Sources

Charles Kaspar, Ph.D., M. Ellin Doyle, Ph.D.
Food Research Institute, University of Wisconsin-Madison

Contents

	Page
Introduction	2
Epidemiology of Non-O157:H7 STEC.....	3
Surveillance and Pathogenicity.....	3
Outbreaks and Sporadic Cases.....	5
Reservoirs of Non-O157:H7	6
Cattle—the primary reservoir	6
Other Ruminants	7
Other Animals.....	8
Transport Hosts.....	9
Routes of Human Infection.....	9
Direct Contact.....	10
Contaminated Food.....	10
Contaminated Water	11
Interventions for Controlling Non-O157:H7 STEC	12
Preharvest Interventions.....	12
Processing Interventions	14
Farm and Zoo Interventions.....	14
Analytical Methods for Detecting Non-O157:H7 STEC.....	14
Introduction.....	14
Enrichment and Isolation	15
Identification of STEC Serotypes	16
Shiga-Toxin Detection	16
Subtyping Methods.....	18
Further Research Needed.....	18
Table 1: Outbreaks of Non-O157:H7 STEC.....	20
Figures.....	22
Reference List.....	23

Introduction

Shiga-toxin producing *E. coli* (STEC) can cause devastating illness, particularly in children, by causing hemolytic uremic syndrome (HUS) leading to kidney failure. Outbreaks of illness caused by STEC have been epidemiologically related to contact with animals and consumption of meat and fresh produce. *E. coli* O157:H7 is the most notorious of the STEC strains causing approximately 73,500 cases in the U.S. each year. CDC estimates that non-O157 STEC are responsible for about 37,000 cases of illness annually but relatively fewer cases of HUS compared to O157:H7. Although many strains of non-O157 STEC appear to be less virulent than *E. coli* O157:H7, a 2008 outbreak of STEC strain O111 in Oklahoma caused illness in at least 314 people, HUS in 17 cases, and one death. (203) Other non-O157 outbreaks in the U.S. have been traced to contaminated lake water, salad greens, and milk.

Numerous *E. coli* strains are capable of producing one or both Shiga-toxins (Stx1 and Stx2) but not all of them are important human pathogens. STEC strains have been divided into 5 seropathotypes: **A**, including the O157 strains that are common causes of outbreaks and HUS in most countries; **B**, non-O157 strains that cause occasional outbreaks but are fairly common isolates from sporadic cases and HUS (examples: O26:H11, O103:H2, O111:NM, O121:H19, O145:NM); **C**, non-O157 strains associated only with sporadic cases; **D**, strains associated with diarrhea, not more severe symptoms; and **E**, strains not associated with human disease. (108;147) Stx2 is the more potent toxin and those strains producing this toxin are generally associated with more acute illness. Other virulence factors are also important, including intimin, adhesions, enterohemolysin, and those involved in type III secretion system that participates in the production of characteristic intestinal lesions. (56;59)

More than 200 virulent non-O157 serotypes have been isolated from outbreaks and sporadic cases of HUS and severe diarrhea in the U.S. and other countries. Serogroups O111, O103, and O26 are among the most frequently detected. (35) The true incidence of non-O157 STEC infections is probably underestimated because standard stool culture methods routinely used in many clinical laboratories do not detect these bacteria. Recently developed analytical methods for STEC strains detect Stx proteins or genes encoding these proteins. (24;113) However, the presence of Stx or its genetic determinants in a sample does not necessarily mean that there are viable STEC bacterial cells.

As with *E. coli* O157:H7, non-O157 STEC serotypes are often associated with cattle and other ruminants and surveys have demonstrated their presence in samples from cattle carcasses, retail beef, and raw milk. (53;130;131;133) Cattle often harbor multiple serotypes, some of which appear to be less of a health risk to humans because they lack one or more important virulence factors. Nevertheless, because microbes can readily exchange genetic information (137), the presence of any STEC in food production environments is of concern. A recent review summarized data on the prevalence of STEC in the beef production chain. (227)

In order to minimize human infections with non-O157 STEC, it is necessary to understand which serotypes are most virulent and all the ways in which people are exposed to these pathogens. A more comprehensive understanding of the epidemiology of infections caused by non-O157 STEC serotypes will lead to improved control methods to prevent illness and reduce economic losses to food producers and processors. This white paper will draw together epidemiological information from the scientific literature and government publications on outbreaks and discuss effectiveness of existing interventions for preventing exposure of humans to pathogenic non-O157 STEC.

Epidemiology of Non-O157:H7 STEC

Surveillance and Pathogenicity

According to published data, non-O157 STEC were first recognized as a possible cause of sporadic cases of HUS in 1975 in France, where hospital records reported that STEC serotype O103 was present in some patients. (148) The earliest reported outbreak, caused by serotype O145:H-, occurred in Japan in 1984. No vehicle of infection was determined for this outbreak. (143) *E. coli* O157:H7 was first identified as a possible human pathogen at about the same time, in a California patient with bloody diarrhea in 1975, and was first associated with a foodborne (ground beef) outbreak of disease in 1982. (228;266)

CDC estimates that about a third of STEC infections in the U.S. are caused by non-O157:H7 serotypes. However, this is likely an underestimate because of the challenges in identification of non-O157 strains. Although there are methods for identification of different serotypes, they are not widely available. In addition, many laboratories do not routinely screen diarrheal stools for shiga toxins and may only attempt to isolate pathogens in cases of bloody diarrhea or if there is a suspected outbreak. (268) It should be noted that there are atypical strains of serogroup O157 designated as O157:H- which can ferment sorbitol and may initially be presumed to be non-O157 strains. Isolates of serotype O157:H- often produce shiga toxins and have been associated with cattle and with severe illness in children. (146;207)

Some surveys in the U.S. and elsewhere indicate that non-O157 STEC may cause diarrhea as frequently as *E. coli* O157:H7 even though they are less commonly identified in cases of severe illness, such as HUS. A 2006 review article (142) reported results from studies in 17 countries indicating that non-O157 serotypes were responsible for 19-100% of STEC infections from which pathogens were isolated. These studies spanned a 10 year period and examined different patient groups (certain ages or geographical areas). So they do not necessarily reflect a greater prevalence of certain serotypes in different countries.

In more recently published surveys, non-O157 serotypes were reported to be significant causes of STEC infections (% = number of non-O157/total STEC identified):

- 80% in a nationwide survey in the Netherlands (256)
- 82% in a laboratory sentinel program in Germany (267)
- 74% in a national surveillance program in Denmark (7;8;35;54;197)
- 13% of HUS cases in France (72)
- 63% in an enhanced surveillance study in Manitoba (252)
- 28% in Ireland in 2008 (93)
- 42-61% in Australia during 2004-2006. (209)
- 24% (2007) and 35% (2008) in Japan (7;8)

Data from summaries of notifiable diseases in the U.S. demonstrate an increasing percentage of cases of STEC infection associated with non-O157:H7 serotypes. In 2002, only 5% of serotyped STEC isolates from human illness were identified as non-O157:H7 strains; in 2005, this had increased to 16% of isolates. (47) Although relative numbers of virulent non-O157:H7 strains may actually be increasing, more frequent testing for shiga toxins and different STEC serogroups undoubtedly explains much of the increase in non-O157:H7 isolates. In Idaho, an enhanced surveillance program targeting a "low" STEC incidence area of the state found that, with more comprehensive laboratory analysis, reported non-O157 STEC incidence increased

from <1 to 11 cases/yr/100,000 population and 56% of serotyped STEC isolates were non-O157 strains. (171)

During 2003-2005, the most common non-O157:H7 strains identified in the U.S. were O26 (19-25%), O103 (14-18%), O111 (13-17%), O45 (5-13%), O121 (6-7%), and O145 (3.4-7.5%).(43-45) Even though a large number of different serogroups are identified in some enhanced surveillance studies of human diarrheal cases, the most common non-O157 STEC strains reported are the six serogroups listed above. Nine to ten serogroups are identified yearly in Wisconsin with the most common serogroups during 2007-2009: O26 (24-32%), O103 (21-34%), O111 (10-29%), O45 (2-17%), and O121 (2-9%). (Data from J. Archer) No cases of HUS have been associated with non-O157 STEC in Wisconsin in the past 3 years, but some strains of all the common serogroups have caused illness severe enough to require hospitalization of some patients. (Data from J. Archer) Virulence factors important for human infection are more commonly carried in these strains resulting in more frequent detection.

Virulence characteristics vary somewhat among STEC strains but all strains, by definition, produce shiga toxin 1 (Stx1) and/or shiga toxin 2 (Stx2). Strains producing Stx2 cause more serious illness than those producing only Stx1. Several variants of these proteins have been described and some are more often detected in certain serotypes or certain host animals. (59) Part of the toxin molecule binds to host cell receptors and facilitates transfer of the toxin into cells. Another toxin subunit has enzymatic activity and interferes with protein synthesis in cells and induces inflammatory responses. In addition to these toxins, virulent STEC often carry the locus of enterocyte effacement (LEE) whose genes code for proteins that form attaching and effacing lesions in the intestines of infected animals. LEE also encodes a type III secretion system to deliver virulence factors to intestinal enterocytes. Many pathogenic STEC also produce enterohemolysin. Several recent reviews discuss these virulence factors in more depth. (21;41;108;139;142)

Some virulence factors may confer advantages to STEC cells in the environment. Shiga toxins are also toxic to *Tetrahymena*, a common fresh-water, protozoan predator of bacteria. STEC may have a survival advantage when these predators are present. (161) In addition to being an important virulence factor in some STEC infections, the serine protease, espP, also aids in attachment to lettuce leaves and may aid survival in the environment. (154;239)

Although non-O157 STEC are generally associated with less severe illness than *E. coli* O157:H7, this may change because bacteria readily exchange genetic material and are constantly gaining or losing genetic information, such as virulence genes. A genomic comparison of virulent STEC strains of serotypes O157, O26, O103, and O111 revealed that they contained a large number of prophages and transmissible integrative genetic elements containing virulence genes. These strains had distinct evolutionary histories and independently acquired the mobile genetic elements coding for virulence factors. (202) STEC strains isolated from patients in the early stages of infection may differ from those isolated from fecal samples several days later, indicating that evolution of virulence characteristics occurs even during a matter of days within a host. (179) Transfer of genes for shiga toxins through bacteriophage transduction could potentially also occur in foods. However, some experiments examining this process in milk, ground beef, salads, and other foods, found that cell numbers would need to be much greater than those normally observed in foods for efficient gene transfer. (137)

Outbreaks and Sporadic Cases

Reports from public health surveillance studies in many (U.S.) states and from other countries indicate that sporadic cases of non-O157 STEC greatly outnumber outbreak cases. (35;127;163;178;197;229) This is also true for *E. coli* O157:H7. According to FoodNet data from 2005, only 23% of 473 confirmed cases of infection with *E. coli* O157:H7 were associated with outbreaks. (48) Approximately 50 non-O157 STEC cases have been identified annually in Wisconsin during 2007-2009 but these were nearly all sporadic cases. (J. Archer)

Outbreaks attributed to non-O157:H7 STEC have been reported from the U.S., Europe, Australia, and Japan. Data on 80 outbreaks, from 1984 to 2009, reported in the literature or government websites are presented in Table 1. It is very likely that other outbreaks have occurred but were not recognized because of the difficulties in identifying and characterizing non-O157:H7 STEC serotypes. Information on some other outbreaks may have been published on foreign language websites or in inaccessible journals and were not included here.

In the U.S., the earliest outbreak occurred in Ohio in 1990 among family members. A 2008 outbreak in Oklahoma, caused by serotype O111:NM, affected 341 patrons and workers at a particular restaurant. Despite an extensive investigation by public health authorities, targeting a variety of foods, food handlers and water sources, no specific source of the *E. coli* O111:NM was identified. (203)

Notable international outbreaks include the 1995 mettwurst outbreak in Australia, caused by serotype O111:NM, which resulted in 23 cases of HUS among 88 persons affected (38) and a more recent outbreak in Norway in 2006 caused by a virulent strain of serotype O103:H25, present in a particular kind of mutton sausage. There were 10 cases of HUS and 1 death among the 18 cases that were recognized after an extensive epidemiological investigation (236). A Danish outbreak in 2007 due to serotype O26:H11, present in a different type of sausage, was much milder with no cases of HUS or death. (75) The O103:H25 strain was reported to produce only shiga toxin 2 while the O26:H11 strain produced only shiga toxin 1.

Another interesting aspect of the Danish outbreak was the epidemiological investigation. Initial interviews with parents of the affected children generated no useful hypotheses. Next, the investigators asked the parents where they shopped in the previous 3 weeks and how much they had spent on food. With this information, they were able to retrieve from the stores' computers exactly what was purchased and they identified one particular organic fermented sausage, bought by several families, as the likely vehicle. The outbreak strain was isolated from the sausage and it was recalled. (75)

Relative importance of different vehicles of infection for outbreaks is depicted in Figures 1 and 2. In 15 outbreaks (18.8%), no vehicle was identified. Person-person contact was reported to be the cause of about 29% of outbreaks and 20% of cases. Many of these occurred in schools and day care situations in Japan. Other outbreaks were traced to meat (9), dairy products (8), water, both drinking water and pool or lake water (8), produce (5), and other food (7). Several small outbreaks (5) occurred among visitors to farms and petting zoos. The "other food" category accounted for about 9% of outbreaks but 27% of cases. This was due to two large outbreaks: the 2008 Oklahoma restaurant outbreak with 341 cases and a 2004 outbreak associated with unpasteurized cider. Compared to outbreak data gathered for a previous white paper on *E. coli* O157:H7 (64), non-O157:H7 STEC strains are much less often associated with meat, water, and produce as outbreak vehicles and much more often attributed to person to person contact or unknown vehicles. (Table 2) These differences are likely due, in part, to the better analytical methods available for *E. coli* O157:H7. *E. coli* O157:H7 is also more virulent than some non-

O157:H7 STEC strains and thus outbreaks are recognized and investigated more rapidly and thoroughly.

Table 2. Comparison of the relative importance of vehicles associated with global outbreaks of non-O157:H7 STEC and *E. coli* O157:H7

Vehicle	non-O157:H7 STEC	<i>E. coli</i> O157:H7(64)
Animal contact	6.2%	9.7%
Water	10.0%	25.6%
Person-person contact	28.8%	6.8%
Dairy	10.0%	12.5%
Meat	11.2%	24.6%
Produce	6.2%	9.2%
Other Food	8.8%	5.8%
Unknown	18.8%	5.8%

Reservoirs of Non-O157:H7 STEC

Understanding the epidemiology of non-O157:H7 STEC serotypes requires a knowledge of where they live and grow in nature (their reservoir) and of how humans come into contact them. Ruminants have been identified as the major reservoir of *E. coli* O157:H7 and also appear to be a reservoir of non-O157:H7 STEC strains. STEC have been isolated from cattle, sheep, goats, and deer. STEC are occasionally isolated from other wild and domestic animals but it is believed that, in many cases, they are present as transients that the animals acquired them from foods or water contaminated by fecal material from ruminants. Nevertheless, some of these transient hosts may be vehicles of infection for humans.

Non-O157:H7 STEC have been detected in numerous species of animals. Two non-O157:H7 STEC outbreaks in Australia were traced to contact with non-ruminants: a 2002 outbreak at a petting zoo with pigs and alpacas infected with STEC serotype O26 (208) and a 2007 outbreak at an animal sanctuary likely caused by koalas and/or kangaroos infected with serotype O55:H80 (110). Both a child with diarrhea and domestic pigeons in Germany were found to harbor the same STEC serotype, O128:H2. (246) and another child in Germany and her cat were found to be excreting identical strains of STEC O145:H- (36). A survey of wildlife meat in Germany found a number of non-O157:H7 STEC serotypes present in deer, wild boar, and wild rabbit meats. Some of these STEC were serotypes that have also been detected in cases of human illness. (182)

Cattle—the primary reservoir

Cattle are probably the most important source of human infections. Of the outbreaks listed in Table 1, 16 appear to be associated directly with cattle. These include two associated with beef, four with “meat,” and six with dairy products from cows. In addition there were four outbreaks associated with contact with animals at farms or petting zoos. Other outbreaks associated with contaminated water and fresh produce may be indirectly associated with cattle.

Wide ranges of prevalences of non-O157:H7 STEC in feces from dairy (0.4-74%) and beef (2.1-70.1%) cattle have been reported from various countries. A total of 193 STEC serotypes were reported from dairy and 261 serotypes from beef cattle. About 12-17% of these serotypes

have also been isolated from cases of human illness. Many of the apparently non-pathogenic strains appeared to be lacking one or more virulence factors. (131;133) While it is difficult to compare results of all these studies because of variations in sampling and detection methods, they do indicate that cattle shed a variety of non-O157:H7 STEC serotypes, some of which are human pathogens.

More recently published studies also demonstrate a large number of serotypes shed by cattle: 26 serotypes detected on organic and conventional dairy farms in MN (51), at least 10 detected on beef carcasses in the Pacific northwest (53) and 31 serogroups detected in dairy cattle in Japan (157). These studies also illustrate an analytical issue that must be considered in analyzing published data. Estimates of prevalence were much higher from PCR or immunoassays detecting shiga toxins or genes coding for these toxins (23-30%) than estimates obtained from isolation of STEC bacteria (6-12%). It is unclear whether the toxin assays are detecting proteins or DNA from viable or dead bacterial cells.

Several studies have reported the STEC are shed more often by cattle during warm months. (227) Most isolates of O26 and O111 in Korean cattle were detected during May-October. (140) Non-O157:H7 STEC were detected more often in Spring and Autumn in Midwestern U.S. beef processing plants. (13) Seasonality may be related to ambient temperatures, age of cattle, and/or type of feed or pasture consumed at different seasons.

A likely source of infection for cattle is feed or water contaminated with feces of other infected animals. Research has documented the survival of STEC O26 for extended periods in manure: up to three months in manure heaps (85) and cow slurry (86) and up to a year in manure-amended soil depending on temperature and soil type (87). Persistence of these pathogens in an environment contaminated with manure is a concern not only for on-farm transmission and reinfection but also for environments such as county fairs and petting zoos where children may be exposed.

Several on-farm studies have documented the acquisition of STEC by calves. Sera and colostrum of dams and sera of newborn calves were found to contain Stx1-specific antibodies which may help protect newborn animals from infection. Antibody titers decreased rapidly during the first 6 weeks and by 8 weeks most calves were shedding STEC. (88) Another study showed that very young calves do not shed STEC (52) and calves shed different STEC strains as they age (238). About 40% of calf infections were estimated to be acquired from other calves (170). Some STEC strains appear to be cleared within a day while other strains persist in calves for several days.(270) Population dynamics of STEC shedding by beef calves from birth to about two years of age was monitored and modeled to understand changes that occurred over time. (70) Results from these studies may suggest new approaches for on-farm control of STEC.

Other ruminants

Sheep have been found to carry a great diversity of STEC but, *E. coli* O157:H7 is infrequently isolated. Non-O157:H7 STEC have been detected in lambs and/or adult sheep from Australia (63), Brazil (259), India (26), Jordan (200), New Zealand (55), Norway (254), Spain (30;204;225), Switzerland (275;277), and the U.S. (135;144). In all of these surveys, multiple STEC strains were detected but O157:H7, if present, was a minor component of those identified. Several serotypes detected in sheep in different countries were similar, indicating that there may be some serotypes that have adapted to colonizing sheep. STEC serotypes that have been associated with human illness were detected in some studies but the majority of STEC strains present in sheep appeared to be of low virulence because they lacked some important virulence factors (intimin, hemolysin, Stx2).

Data gathered at a commercial lamb processing plant in the U.S. revealed that the prevalences of *E. coli* O157:H7 on pelts, preevisceration carcasses, and postintervention carcasses were 12.8%, 1.6%, and 2.9% respectively. For non-O157:H7 serotypes, prevalences were 86.2%, 78.6%, and 81.6%, respectively. A total of 69 different non-O157:H7 serotypes were identified. About 4% of these serotypes have previously been associated with severe human illness. (144)

Goats are another reservoir of STEC. Of 13 caprine dairy herds surveyed in Ireland, STEC were isolated from milk filters from 3 farms. Serotypes O157 and O26 were each detected twice. (193) A longitudinal study of two dairy goat farms in Spain documented chronic shedding of STEC by many adults on both farms but more sporadic shedding by kids. On one farm kids were carrying STEC bacteria within 1 week of age while on the other farm, it was four months before shedding began. Fewer adults were frequent shedders on the latter farm. Serotypes identified were primarily O33, O76, O126, O146, and O166. None of the isolates produced intimin and they appear to be adapted for colonizing the goat intestine. (205) Non-O157:H7 STEC have also been detected in goats in Jordan (200;249), Bangladesh (138), and Vietnam (128).

Buffaloes. Nearly one quarter of the buffalo slaughtered at a facility in Bangladesh contained non-O157:H7 STEC of several serogroups. Most strains produced Stx1 but many did not produce hemolysin or some other virulence factors. *E. coli* O157:H7 was isolated from 14.4% of the buffalo. (138) A survey of 98 farms in central Vietnam revealed that 70% of farms had STEC-positive buffalo. However a minority (5%) of the serotypes identified were commonly associated with human illness. (128)

Guanaco (*Lama guanicoe*). STEC O26:H11 was isolated from a two-month old guanaco with severe diarrhea in Argentina. (181)

Deer and elk are present in significant numbers in some environments also used by cattle, sheep and goats and their droppings may contaminate fresh produce in the field and surface waters. Cultures from fecal pellets from native Idaho ungulates revealed that about 19% were positive for Stx which is about the same prevalence reported for cattle in that state. (80;96) There have been numerous reports of *E. coli* O157:H7 in wild deer in the U.S. and other countries (119) and a few cases of human illness attributed to deer meat (152). Free-ranging wild sheep and deer in Spain were found to shed at least 11 non-O157:H7 STEC serotypes with O146 being the most commonly detected serogroup. (230) Several captive ruminants in an Argentine zoo, including alpaca, antelope, deer, eland, sheep, and yak, shed non-O157:H7 STEC. (168;222)

Other animals

Swine have been found, in several studies, to be infected with both *E. coli* O157:H7 and non-O157:H7 STEC strains.(22;82;120;141;150;151;199;235;276) while in other surveys, only non-O157:H7 STEC were detected (155;264). Although virulent STEC strains have been found in a few samples from swine, most researchers conclude that swine are not an important source for human infections because the STEC strains isolated from these animals often lack some important virulence genes and differ from strains usually isolated from cases of human illness. (62;155;264;276)

Horses do not appear to be an important reservoir for STEC. Only one of 400 fecal samples from horses in Germany tested positive for STEC (serotype O113:H21) and one of 100 horse meat samples tested was positive for STEC (serotype O87:H16). (218)

Rabbits, both wild individuals and animals being raised commercially on farms, were reported to harbor non-O157:H7 enterohemorrhagic *E. coli*. (91;164;231)

Poultry have occasionally tested positive for *E. coli* O157:H7 (64) but so far there are no reports of non-O157:H7 STEC in poultry.

A **Cat** and a 2-year old German girl with bloody diarrhea were found to excrete the same STEC serotype O145:H-. The cat had no symptoms but was found to excrete this STEC strain for several months and was apparently the source of the child's original infection and/or reinfection. (36)

Dogs in Brazil were reported to harbor non-O157:H7 STEC that caused diarrhea. (58)

Shellfish in contaminated waters are known to concentrate some pathogens such as *Cryptosporidium*. *E. coli* is present in human sewage and the possibility exists that pathogenic strains such as STEC could be present in lakes, rivers or coastal waters contaminated by sanitary sewer overflow or runoff from fields containing fecal matter from domestic or wild animals. There have been reports of non-O157:H7 STEC detected in shellfish collected from coastal areas of France (100) and India (160;175). However, it appears that STEC strains are not a significant contaminant of shellfish.

Transport Hosts

Birds are a potential transport host for STEC because some wild birds harbor these bacteria and might spread them around a farm environment. There are a number of reports of non-O157:H7 STEC in both captive (ornamental, racing) pigeons (78;102;246) and in feral pigeons in the city or countryside (102;156;190). Stx2f originally described from pigeon isolates (234) has been detected in a number of human diarrheal STEC isolates. (221) Although a possible case of domestic pigeon-to-human transmission of non-O157:H7 as been reported (246), a study in Colorado indicated that wild pigeons may not be a major route of transmission of STEC. (214)

Other wild birds may also carry non-O157:H7 STEC. A starling on a Danish farm was found to harbor STEC serotype O2:H29. (198) STEC serotype O20 was detected in an Oriental turtle dove and serotype O147 was detected in a barn swallow living near Tokyo Bay. (156) Stx2 was detected in feces of 30 wild bird species (of 99 species tested) in the UK but strains were not serotyped. (125)

Rodents, including mice and rats, are also potential transport hosts for STEC. However, STEC have not been frequently reported from these animals. There is one report of STEC serotype O136:H12 from a rat on a cattle farm in Denmark. (198) Among animals tested at an Argentine zoo, STEC serotype O146:H28 was detected in a cavy (related to guinea pigs). (168)

Flies and beetles, collected on farms with animals shedding *E. coli* O157:H7, have been found to contain detectable levels of these bacteria. (64) These insects frequent fecal deposits and may transfer these pathogens to foods, feed and water. STEC were recently isolated from flies collected at pig pens and in cattle barns but were not serotyped. (90)

Routes of Human Infection

Ruminant fecal material is believed to be the ultimate source of a large percentage of human non-O157 STEC infections. A study in Germany found that there was a positive association between illness caused by a number of non-O157 STEC serotypes and the density of cattle in an area. From data on over 3000 STEC cases, analyses indicated that risk for infection increased by 68% /100 additional cattle/km². Increased risk varied for different serotypes but was greatest for O111. (81) A similar association has been documented for *E. coli* O157:H7. (255) Sporadic STEC cases have been traced to contact with cattle on farms. (118;196)

Fecal material may contaminate meat during slaughter, may be washed or blown into lakes or drinking water sources, or may be deposited on fruits and vegetables by use of manure for fertilization or sewage-contaminated water for irrigation. Some animals, such as insects, birds, and rodents, may transport these bacteria from feces to drinking water or foods. In addition, non-O157 STEC may be inadvertently ingested by persons interacting or working with animals. Humans may therefore acquire infections through direct contact with an infected person or animal or their environment or through food, drinking water or surface water containing STEC contaminated fecal material from an animal or human. (115)

Direct contact.

Numerous outbreaks of enteric zoonotic disease have been associated with animal exhibits at fairs, zoos and other venues. *Cryptosporidium*, *Salmonella* spp., and STEC are the pathogens most commonly identified in these outbreaks. (119;167) Several non-O157 STEC outbreaks among children who visited farms (1;245) or petting zoos (110;208) resulted from direct contact with animals and their environment followed by inadequate hand washing. *E. coli* O26:H11 and O111:H- can survive in cattle feces for 10-12 weeks at 15°C (89) and STEC may persist on surfaces at farms and zoos for extended periods (even after animals have stopped shedding) if there is sufficient moisture and temperatures are not excessive. (119) An outbreak of non-O157 STEC at a Minnesota farm day camp occurred in two consecutive years despite attempts to clean the premises and encourage hand washing. (245)

Contact with domestic animals has also been a route of STEC infection. A cat and a 2-year old German girl with bloody diarrhea were both found to excrete O145:H-. Although the cat had no symptoms, it excreted this strain for several months and was apparently the source of the child's original infection and/or reinfection. (36) Another child with diarrhea and some pigeons harbored the same STEC O128 strain. (246) Pet rabbits have also been reported to harbor non-O157:H7 enterohemorrhagic *E. coli*. (91)

Person-to-person spread of non-O157 STEC has been the primary mode of infection in outbreaks in day cares, schools and senior care facilities. (7;8;35;54) In many other outbreaks, some cases who consumed contaminated food or water, passed the infection directly to friends or others in their family. Although most people apparently stop shedding STEC bacteria within a week or so of recovering from illness, there are some people who continue to shed bacteria for weeks or months afterwards. A study in a German sausage factory over a 21 month period demonstrated that one healthy worker excreted non-O157 STEC intermittently for 7 weeks while another symptomless worker excreted STEC for nearly ten months. (92)

Contaminated food

Beef, lamb, and mutton can be contaminated during slaughter and processing by exposure to feces or hides containing non-O157 STEC. A 2007 review stated that reported levels of non-O157 STEC in whole cattle carcasses, ground beef, retail beef cuts, and sausage were 1.7-58%, 2.4-30%, 11.4-49.6%, and 17-49.2%, respectively. (129) Beef trim, which is ground to make hamburger, is believed to be an important source of STEC contamination in ground beef. A survey of boneless beef trim from Australia (220 samples), New Zealand (223 samples), Uruguay (256 samples), and the U.S. (487 samples) revealed that non-O157 STEC were present in 10% of the New Zealand samples and in about 30% of the other samples. (31) Other surveys have reported the STEC prevalence to be: 15% in ground meat in France (216), 1.5% in beef in Japan (113), 40% in ground lamb in Australia (14), 1% in horse meat in Germany (218), and

24% in buffalo meat in India (116) Procedures for collecting samples and performing analyses differed among these studies, so results are not directly comparable.

Milk from dairy cows, sheep, and goats may be contaminated with *E. coli* and other bacteria from the environment. A review in 2005 summarized numerous surveys that detected STEC in milk and dairy products. Contamination with *E. coli* is generally low and some of the STEC strains detected in raw milk have not been associated with human disease. (133) Some non-O157 STEC may be more prevalent in milk than *E. coli* O157:H7. STEC strains have been shown to survive various steps in cheese-making so that raw milk cheeses are potentially a vehicle for STEC infections. (16)

Most later surveys also report a relatively low prevalence of non-O157 STEC in raw milk but there were some positive samples indicating the need for proper treatment of milk. (173;174) STEC strains were isolated from 11% of bulk milk tank samples, 4% of cheese curds, and 5% of cheese in Spain (226) and in 21% of raw milk samples in France (216). In a Swiss study, non-O157 STEC were isolated from 16 of 744 raw milk hard and semi-hard cheeses. (248) Analyses of 40 STEC strains isolated from raw milk and cheese in France, found that most of the strains lacked some virulence factors found in isolates from human disease. (220) Proper pasteurization kills *E. coli* so outbreaks of STEC due to contaminated dairy products are usually associated with unpasteurized milk. (2;4;61) But there has been an outbreak due to post-pasteurization contamination. (189)

Field and greenhouse experiments have demonstrated that both *E. coli* O157:H7-contaminated manure and irrigation water may cause contamination of vegetables and this is probably true for non-O157 STEC as well. Manure is a valuable fertilizer for crops but manure containing STEC may be a source of contamination for vegetables or fruits that are not normally cooked before eating. In one study, non-O157 STEC were able to survive for 42 days in manure heaps that were turned and for 90 days in unturned heaps. (85) STEC O26 survived for at least 90 days in cow slurry (an effluent comprised of feces, urine, water, spilt feed, and bedding) (86). Serotype O26 was also detectable in manure-amended loam soil for more than 9 months at 4°C and for more than 6 months at 20°C (87).

Foods can also be contaminated with STEC by cross-contamination during food preparation and by infected workers who don't practice good hygiene. There have been several outbreaks attributed to restaurant food. Cross-contamination in food preparation areas or infected food handlers might have contributed to these outbreaks. An outbreak in a prison was traced to a food handler. (46)

Contaminated water

Water used for drinking or recreation has been reported as the vehicle of infection for 7 outbreaks. One outbreak in 1988 in Czechoslovakia was associated with tap water. Several outbreaks occurred among children playing or swimming in pool or lake water. Other infected children may have been the source of bacteria for these cases. Other outbreaks were traced to water consumed at summer camps. Fecal material from domestic and/or wild ruminant animals may have contaminated lakes, rivers, and some "drinking water."

Surveys of some surface waters have detected *stx* genes in beach and stream water in a park in Pennsylvania (243), and in river water in Michigan and Indiana (67) and in India (223). The significance of these findings is unclear because the presence of *stx* genes was not correlated with numbers of viable bacteria present. Some strains of the non-O157 STEC serotypes O26 and O111 have been reported to survive in untreated well water for over 56 days at 10°C. Cells die off more quickly at 22°C but do persist in significant numbers for four weeks. (265)

Interventions for Control of Non-O157:H7 STEC

Research on prevention of STEC contamination of foods and water and strategies to kill or severely limit growth of any STEC that might be present in foods has concentrated primarily on *E. coli* O157:H7. In the following sections, data will be presented for non-O157 STEC when available and summarized for *E. coli* O157:H7 or other *E. coli*. Susceptibility of non-O157 STEC to various intervention techniques is probably similar to that of other *E. coli* although there are known differences among strains in acid tolerance and sensitivity to some other agents. Some recent reviews discussing intervention techniques have been published. (119;134;224;257)

Effects of processing technologies on *E. coli* and other bacteria in meat were recently reviewed. (10) Among the procedures discussed were: irradiation, high hydrostatic pressure, natural antimicrobials, active packaging, and thermal treatments. Comparisons between *E. coli* O157:H7 and non-O157 STEC were not discussed but conditions generally effective against *E. coli* were described.

Pre-Harvest Interventions

Dietary Interventions

Feed. Results of experiments published about ten years ago indicated that different diets fed to cattle may affect concentrations of *E. coli* O157:H7 shed in feces. Feeding of high grain rations to feedlot cattle to increase feed efficiency causes some starch from the grains to escape fermentation in the rumen and pass to the hindgut where it is fermented by other bacteria. This can change the pH of the rumen and hindgut thereby affecting survival of some bacteria, such as STEC strains. A recent review paper discussed various experiments conducted since that time to determine whether dietary interventions could reduce numbers of STEC bacteria excreted by cattle. (37) Subsequent experiments have shown that dietary differences do affect *E. coli* populations in cattle but the effects varied in magnitude and impact. Some studies suggest that the tannins and phenolic acids in forage may be the important components affecting shedding of STEC while other experiments, in which barley and distillers grains were fed to cattle, demonstrated an increase in shedding of *E. coli* O157:H7. Dietary experiments that tested for genes coding STEC virulence factors in cattle feces (which would measure effects on all STEC not just *E. coli* O157:H7) suggested that a diet with more roughage may reduce concentrations of STEC bacteria that cause human disease. However, the cattle in this study were excreting low levels of STEC and further experiments are needed to determine the significance of these results. (95)

Probiotics, commensal bacteria fed to animals to reduce numbers of pathogens, have been suggested as a strategy to prevent growth of STEC in young ruminants. In one experiment with calves, a three strain mixture of non-pathogenic *E. coli* were fed to calves three days after challenge with one of three STEC serotypes. The probiotic *E. coli* mixture decreased fecal shedding of O157:H7 and O111:NM but did not affect shedding of O26:H11. At necropsy, all of the calves challenged with O26:H11 and four of twelve calves challenged with the other STEC still harbored viable STEC. (253) *E. coli* strains used as probiotics may produce colicins that kill

STEC strains. (232) Other experiments have tested the effects of lactic acid bacteria on STEC survival. Some inhibition of STEC has been observed in tests in vitro. (98;126) Some variability in practical results of tests with probiotics may be due to different management practices including the effects of subtherapeutic levels of antibiotics fed to livestock to enhance growth. (119)

Bacteriophages are viruses that can kill bacteria and have been proposed as potential control agents for STEC. In experiments with an artificial rumen system, phage D22, specific for *E. coli* O157:H7, eliminated these bacteria from the fermentor within four hours. However, in experiments with lambs, inoculated first with *E. coli* O157:H7 and then two days later with the phage, there was no decrease in numbers of *E. coli* O157:H7 shed. The phage did not persist in these animals. (11) To avoid the problem of inactivation of phage in an animal's digestive tract, another experiment tested the effects of two phages (one of which could also kill some non-O157 STEC strains), the phages and *E. coli* O157:H7 were applied directly to the rectoanal junction of steers. Phage therapy reduced the average number of *E. coli* O157:H7 detected in feces but did not eliminate these bacteria from most of the animals. (240) Although bacteriophages have not yet emerged as a practical preharvest solution to STEC shedding by ruminants, some research continues to find other phages that might be more effective. (262)

Chlorate is metabolized by some bacteria, including *E. coli*, to a toxic compound, chlorite. Feeding chlorate to cattle in feed and water prior to slaughter can significantly reduce concentrations of *E. coli* O157:H7 and other *E. coli* in feces at slaughter. This is a potentially useful strategy for reducing contamination of meat during processing at slaughter facilities. (6) Chlorate in drinking water is also effective in reducing *E. coli* populations. (119)

Drinking Water may be contaminated by fecal material and is known to be a source of infection for cattle. (241) Four chemical treatments of drinking water using lactic acid, acidic calcium sulfate and one of the following: benzoate, caprylic acid, butyric acid, chlorine dioxide resulted in >3 log reduction in numbers of *E. coli* O157:H7, O26:H11, and O111:NM in contaminated trough water. However, cattle consumed much less water when these chemicals were present, so they should not be used continuously. (274)

Vaccines

Some infected calves develop an immune response to STEC and vaccines targeting some important STEC proteins may be useful in preventing the establishment of STEC in calves. A recent review article mentioned 7 vaccines that have been described in the literature. There is not much information available on most of them because of proprietary considerations. Some have been tested in cattle but others have apparently been tested only in pigs or laboratory animals. Some vaccines, if demonstrated to be effective in cattle, may offer protection against non-O157 strains if they induce antibodies to a common virulence factor. (257) Two recent articles described some new vaccines but they have not been tested in cattle as yet. (105;169) Large-scale testing of a vaccine to reduce carriage of *E. coli* O157:H7 in cattle has begun in Colorado. A New York Times article on December 4, 2009 described the program and the obstacles faced by vaccine producers in getting approval for use of this product.

Processing Interventions

Many interventions that aid in control of *E. coli* O157:H7 are likely to be effective for non-O157 STEC also. Effectiveness of interventions for decontaminating meat were recently reviewed. (10) Use of hot water and lactic acid washes and steam to clean carcasses effectively reduce contamination with *E. coli* O157:H7. (32;158) Intervention techniques to remove *E. coli* O157:H7 from the surface of beef carcasses were found to be similarly effective against O26:H11 and O111:H8. (57)

Stress tolerance to heat, salt and acid has been observed in many STEC strains and should be considered when devising interventions in food processing. Some non-O157 STEC are more susceptible or resistant to stresses so that effectiveness of procedures needs to be tested with more than one serotype. For example, *E. coli* O157:H7 was found to be more resistant to acid in a model stomach system than some non-O157 STEC serotypes. (18) However, other tests in different media found *E. coli* O157:H7 to be more sensitive to acid than other non-O157 STEC. (20;162;186) Acid resistant non-O157 STEC survive longer in fermented raw sausage than non-acid resistant strains. (188) Both sodium lactate and sodium benzoate inhibit non-O157 STEC but the extent of inhibition is temperature dependent. (117;166)

STEC serotypes do not grow at refrigeration temperatures but can remain viable in food for extended periods in the cold (211) and *E. coli* O157:H7 can survive at least for several days in meat and yogurt when frozen at -18°C (104). Thermal treatments can destroy STEC but one study reported that STEC O26 was less heat sensitive than *E. coli* O157:H7 in minced beef heated at 55°C. (66)

Effectiveness of sanitizers and disinfectants have been tested against STEC O111 and O26. (159;242;258) Several STEC serotypes tested were more resistant to desiccation on paper discs than *Shigella* and non-pathogenic *E. coli*. STEC also survived for months in chocolate. (17;121)

Zoo and Farm Environmental Interventions

CDC published in 2009 an updated version of recommended measures to prevent disease associated with animals in public settings. Million of human-animal interactions occur annually in a variety of settings. Hand washing is the most important preventive step for reducing disease transmission. In addition, CDC recommends prohibition of food in animal areas, education of visitors about disease risk and prevention, proper care and management of animals, and transition areas between animal and non-animal locations. The updated guidelines also discuss risks associated with baby poultry, reptiles, rodents, and aquatic animals. (49)

Composting of manure from animals shedding STEC can eliminate STEC when temperatures and the presence of other bacteria are optimized. (97)

Analytical Methods for Detecting Non-O157:H7 STEC

Introduction

E. coli O157:H7 can usually be readily identified in the laboratory because of its inability to ferment sorbitol or cleave the fluorogenic substrate 4-methylumbelliferyl-B-d-glucuronide within 24 hours which distinguishes it from other *E. coli* and most of the other bacteria in its

environment. Nearly all *E. coli* O157:H7 produce shiga-like toxins or harbor genes (*stx*) encoding the toxins so a culture-positive result is assumed to be positive for STEC. It should be noted that there are atypical strains of serogroup O157, designated as O157:H-, which can ferment sorbitol and may initially be presumed to be non-O157 strains. Isolates of serotype O157:H- often produce shiga toxins and have been associated with cattle and with severe illness in children. (146;207)

Detection and identification of non-O157:H7 STEC serotypes in a timely fashion is more difficult. These strains do ferment sorbitol so they are not detectable on sorbitol MacConkey agar plates. Even though six non-O157 serogroups (O26, O45, O103, O111, O121, O145) cause most of the reported cases of non-O157 infection, over 150 STEC serotypes have been associated with illness. In addition, not all strains of these serogroups produce *stx*. Therefore, CDC recommended in a recent report that laboratories simultaneously (1) test samples for the toxins with enzyme immunoassays or for *stx* with PCR methods and also (2) isolate and grow the bacteria in pure culture. (99) If *stx* is detected, then cultures will be immediately available for serotyping and molecular characterization.

Over 100 reports in scientific journals describe analytical methods for detection of non-O157 STEC. Immunoassays, PCR methods, and molecular analytical methods, developed in the past five years, are highlighted in the discussions below. Some useful descriptions of culture and enrichment methods are also included. Several comparisons of the accuracy of different methods and evaluations of commercially available detection methods will be discussed in the final section along with future research needs.

Enrichment and Culture

Isolation of non-O157:H7 serotypes from animals and foods containing large numbers of a variety of other bacteria is challenging because of the genetic and biochemical diversity of these STEC and their similarity to some non-pathogenic bacteria. In a 2006 review article, enrichment/culture protocols, described in 132 papers published since 1997, were discussed. (261) Many researchers used media containing bile salts to inhibit non-Enterobacteriaceae and antibiotics, such as novobiocin which inhibit primarily Gram-positive bacteria. Incubation times and temperatures varied. However, because of variations in experimental procedures, no definitive conclusions could be drawn about the relative effectiveness of the protocols for enrichment from different environmental samples.

Other antibiotics (cefixime, vancomycin) and selective agents (tellurite) have been used to inhibit non-STEC bacteria (132) but there are reports that some non-O157:H7 STEC are sensitive to these agents. (15;206) Other experiments demonstrated that growth of some non-O157:H7 STEC is inhibited by novobiocin in media and use of this antibiotic was not recommended. (260) Universal preenrichment broth (which does not contain antibiotics), incubated at 42°C, was reported to be more effective for detection of STEC O26 and O157 on beef, poultry, and radish sprouts than modified *Escherichia coli* broth with novobiocin. (145)

An acid enrichment procedure was recently reported to substantially decrease background flora in fecal specimens and enhance recovery of STEC strains. (124) In a comparison of three enrichment protocols, a USDA procedure, an FDA procedure and acid enrichment, *stx2* gene was detected in more samples of swine waste treated with acid than samples enriched by the other methods. All strains were non-O157:H7 STEC. The acid enrichment media was found to support the growth of all of 31 STEC strains tested while FDA media supported growth of 23 strains and USDA medium supported growth of 5 strains. (101)

Identification of STEC Serotypes

Culture methods

There are no completely reliable culture methods for identifying non-O157 STEC strains, although research indicates that some specific nutritional requirements or capabilities are associated with certain STEC serotypes. O26 strains that produce shiga toxins are often unable to ferment rhamnose and can usually (but not always) be distinguished from non-toxicogenic strains on rhamnose MacConkey agar. (77) A procedure using a consecutive series of differential and confirmation media was developed to distinguish 4 STEC serotypes (O26, O103, O111, O145). Samples of artificially contaminated food and fecal samples were first enriched in media containing novobiocin, vancomycin, rifampicin, bile salts and tellurite. On the first differential media containing sorbose and sucrose, the non-O157 STEC strains produced different colored colonies. These were then plated to media containing d-arabinose, d-raffinose, l-rhamnose, or dulcitol for confirmation. Isolation efficiency of all serotypes from different sources was 100%, 82.3%, 88.5%, 65.9%, 64.3%, and 13.6% for STEC in raw milk, cheese made from pasteurized milk, cheese made from raw milk, ground beef, fermented meat, and cattle feces, respectively. (219)

Immunoassays

E. coli serotypes can be identified by immunoassays that target the O and H antigens on cell surfaces. Antisera for the most common non-O157 serotypes (O26, O45, O103, O111, O121, O145) are available commercially and can be used to identify many STEC isolates. (99) Antibodies may be coated on immunomagnetic-separation beads and are potentially useful for detecting cattle shedding a large number of STEC of certain serotypes. But these methods are not as reliable if STEC numbers are low. (109) Immunomagnetic beads have also been used to detect O26 and O111 in ground beef. (201)

PCR for serotype specific genes

PCR (polymerase chain reaction) methods targeting DNA variants specific to different serotypes have been developed recently. These are often combined with PCR assays detecting genes coding for shiga toxins or other virulence markers. Some assays target a gene associated with the O antigen such as the *wzx* gene in serotype O26 (172) and in serotype O103 (215) while others target genes associated with both the O and H antigens as in methods described for O111:H8 (69) and O26:H11 (68). More recent multiplex methods detect specific genes present in O-antigen gene clusters of four or five different O groups. (23;83;84;187;217) These procedures can identify serotypes isolated from foods and fecal samples.

Shiga-Toxin Detection

Immunoassays for toxins

Six commercial immunoassays have been approved by FDA for the diagnosis of STEC infections (99):

- Biostar OIA SHIGATOX, an optical immunoassay which does not distinguish between stx1 and stx2, can detect toxin in broths and fecal samples. (251) (This will be withdrawn from market in 2009.)
- Duopath Verotoxin test detects and differentiates shiga toxins 1 and 2 in less than one hour in cultures of isolated cells. (212)

- Immunocard STAT!EHEC also differentiates shiga toxins 1 and 2 in less than an hour in enrichment broths and cultures of isolated cells. (99)
- Premier EHEC does not distinguish between Stx1 and Stx2 and takes several hours to perform. It can detect toxins in stool samples and enrichment broths. Stx concentrations in fecal samples are typically very low, however, and detection is better in enrichment broths. (153)
- ProSpecT Shiga Toxin *E. coli* Microplate Assay does not differentiate Stx1 and Stx2 and takes several hours to perform. It can detect toxins in stool samples as well as enrichment broths. Results are generally better in testing broth cultures.(94)
- VTEC Screen "Seiken" does differentiate Stx1 and Stx2 and takes several hours to perform. It detects toxins in cultures of isolated cells. (42)

An evaluation of the Ridascreen Verotoxin Immunoassay published in 2007 noted that it could detect all known variants of Stx1 and Stx2 in routine screening of bacterial isolates. (24) A more recent evaluation of this assay and the Premier EHEC and ProSpecT assays found that none of the tests could detect some Stx2 variants. The ProSpecT assay was about ten-fold less sensitive than the other two assays. Premier EHEC assay may be useful in screening cattle. (271)

Nanoparticle assay for toxin proteins

Stx bind to cell surface receptors containing terminal Gal- α 1,4-Gal disaccharides. Glycopolymers, with Gal- α 1,4-Gal disaccharides on their surfaces, were found to change color from purple to brown when bound to stx. These nanoparticles could distinguish between *E. coli* strains that did and did not produce Stx within 10 min.(194)

PCR for toxin genes

PCR assays for *stx* genes are generally designed for testing isolated cells from media or bacteria growing in enrichment broths rather than bacterial DNA extracted directly from foods or fecal specimens. PCR procedures have been developed and evaluated for identification of STEC from human stool samples (103) cattle feces (237), and foods (9;84). Some multiplex PCR assays are designed to screen for different types of diarrheagenic *E. coli* targeting virulence genes found in enterohemorrhagic, enteroinvasive, enteropathogenic, enterotoxigenic, and enteroaggregative strains. (107) An evaluation of the GeneDisc assay, a multiplex assay targeting genes for *stx*, intimin, and DNA sequences characteristic of O26, O103, O111, and O157, found that it was very sensitive and capable of detecting 2-3 STEC colonies in a lawn of 50,000 bacteria. (25) A highly sensitive immuno-PCR utilizing an immunoassay with antibody capture and DNA amplification detected as little as 10 pg purified *stx*2/ml (compared to 1 ng detected by a commercial immunoassay. (273)

LAMP (loop mediated isothermal amplification) assay for toxin genes

This method for nucleic acid amplification differs from PCR in that 4 or 6 primers are used to amplify the target gene at a single temperature step. Amplification products can be detected by turbidity because a by-product of the reactions, magnesium pyrophosphate is insoluble. Turbidity caused by this precipitate correlates with the amount of DNA synthesized. (112) DNA was extracted from an enrichment from ground beef and tested with a LAMP assay targeting *stx*. Several STEC serotypes were detected. (111)

Comparison of different stx detection methods

STEC strains have been detected in the past by determining their cytotoxic effects on Vero (monkey) cells in culture. PCR assays for stx give results that are more than 90% in concordance with the Vero cell assay.(272)

Subtyping Methods

Particularly in suspected outbreak situations, it is important to specifically identify the causative pathogen in order to trace pathways of contamination and determine the extent of outbreaks. There is great genetic diversity within STEC due to insertions and deletions in certain parts of the chromosome and genetic information carried by bacteriophages. (202) PFGE (pulsed field gel electrophoresis) is a widely used technique that analyses patterns of chromosome fragments generated by restriction enzymes that cause breaks at certain DNA sequences. Such analyses aided in detection of foods associated with an outbreak caused by STEC O103:H25 in fermented sausage in Norway in 2006 (236) and in tracing the contamination of ice cream associated with a Belgian outbreak(60). Other molecular analyses of multiple genetic loci have provided information on important virulence mechanisms and evolutionary relationships among various STEC strains. (56;180;202;244;250)

Future Research Needs

More recently published studies also demonstrate that a large number of *E. coli* serotypes may be present in animals: at least 10 detected on beef carcasses in the Pacific northwest (53) and 31 serogroups detected in dairy cattle in Japan (157). Estimates of prevalence are much higher from PCR or immunoassays detecting shiga toxins or genes coding for these toxins (23-30%) than estimates obtained from isolation of STEC bacteria (6-12%). (157) Many of these STEC strains may not be virulent but it would be useful to know how readily these strains can exchange genetic information and acquire virulence factors. Although multiplex PCR methods are available for confirmation of virulence genes in isolates, initial screening methods that segregate the diverse populations of *E. coli*, that are commonly encountered in clinical, environmental, and food samples, would facilitate detection and identification of seropathotypes. A combination of effective selective and differential plating media and molecular typing methods are needed to make accurate and comparable prevalence determinations and environmentally track specific serotypes or strains.

Many studies have documented the effects of various intervention techniques on *E. coli* O157:H7. Some interventions should also be tested on pathogenic non-O157 serotypes. The need for these validations arises from the variability of growth and survival properties within a given species such as *E. coli*. With multiple serotypes comprising the non-O157 STEC, there is greater diversity and a need to ascertain if certain serotypes, seropathotypes, or strains have growth or survival properties that differ significantly from *E. coli* O157:H7. Because of variations in resistance to environmental stresses, it is possible that the lethality of interventions and processes used to control *E. coli* O157:H7 will need to be evaluated with a select set of non-O157 STEC.

To more completely define the epidemiology of non-O157 STEC, additional information on animal, environmental, and asymptomatic human hosts is needed. A more complete understanding of localization within hosts as well as the growth, persistence and dissemination in the environment would be beneficial. As with *E. coli* O157:H7, identifying where the pathogen is located in and on the animal can lead to more effective harvesting and processing strategies to reduce contamination of raw food products. Identification of preferred environmental niches can

lead to possible on-farm interventions and reduction of pathogen prevalence. Lastly, the incidence of person-to-person transmission by non-O157 STEC requires further investigation to fully understand the role humans may play in the dissemination of this diverse group of pathogens.

Table 1. Reported Outbreaks Caused by Non-O157 STEC.

Date	Strain	Location	Cases	HUS	Deaths	Vehicle	Ref	Other details
1984	O145:H-	Japan	100		0	Unknown	(143)	school
1986	O111:H-	Japan	22	1	1	Unknown	(12)	orphanage
1988	O111:H2	Australia	2	2	0	Unknown	(12)	family
1988	O26:H11	Czech Repub.	5	5	1	Water, tap	(27)	
1990	O111:NM	US: OH	5	1	0	Unknown	(12)	family
1991	O111:H-	Japan	234			Unknown	(12)	primary school
1992	O111:B4	France	26	10	0	Person-person	(33;39)	school
1992	O111	Italy	9	9	1	Unknown	(4;40)	
1992-93	O119:B14	France	4	4	1	Cheese, unpasteurized	(4;61)	bovine and caprine milk
1994	O103	France	4	4	0	Cheese, raw caprine milk	(4)	
1994	O104:H21	US: MT	18	0	0	Milk, pasteurized	(79;189)	post-past. contamination
1995	O111:H-	Spain	13			Water, drinking	(29)	summer camp
1995	O111:NM	Australia	88	23	1	sausage	(38;71;213)	mettwurst
1995-96	O157:H-	Germany	28	28	3	Sausage	(5)	
1996	O118:H2	Japan	131	0	0	salads	(114)	junior high school
1997	O026:H11	Japan	27	0	0	watermelon, sprouts, spinach	(122)	Nursery
1999	O026	Ireland	10	0	0	Unknown	(28)	child care facility
1999	O026:H11	Germany	3	3	0	Unknown	(184)	3 children, small town
1999	O111:H8	US: TX	58	2	0	Unknown: salad and/or ice?	(19;34)	teenage campers
1999	O121:H19	US: CT	11	3	0	water, lake	(177)	swimming in lake
2000	O026:H11	Germany	11	0	0	beef	(269)	day care centers
2000	O103	US: WA	18	+		Punch	(35)	
2000	O111	US: UT	102			Water, drinking	(165)	also O157:H7 & <i>C. jejuni</i>
2000-01	O051:H11, O111	US: MN	7	0	0	Calves, contact	(245)	farm day camp, 2 years; other enterics children
2001	O026:H11	Japan	11	0	0	water, river	(123)	water used for washing
2001	O26	US: MN	4	0	0	water, lake	(35)	
2001	O26:H-	Austria	2	2	0	Milk, cow's, raw	(2)	
2001	O111	US: SD	3	0	0	Person-person	(35)	day care centers
2001	O111	US: MN	2	2	1	Unknown	(183)	
2002	O148	France	10	2	0	Mutton, lightly roasted	(73)	wedding reception
2002	O26	Australia	5			Animal contact: pigs, alpacas	(208)	petting zoo
2002	O157:H-	Germany	38	38	4	Unknown	(3)	
2003	O111	Australia	13	0	0	Person-person	(54)	old folks home

2004	O086:H27	Australia	4	2	0	Cattle, person-person	(191)	1st case from cattle farm,
2004	O111:NM	Japan	107	0	0	meat	(149)	students traveling to Korea
2004	O121: H19, O121:H7	Japan	63	0	0	Animal exposure: cattle	(1)	children, farm visit
2004	O111	US: NY	213			Cider, unpasteurized	(263)	also <i>Cryptosporidium</i>
2005	O26	Italy	6	6	1	Milk products, buffalo	(173)	
2005	O026 & O80	France	16	16		cheese from unpasteurized cows' milk	(72)	
2005	O026:H11	Japan	12	0	0	Person-person	(136)	kindergarten, also norovirus
2005	O45	US: NY	52	0	0	Food handler	(46)	prison
2006	O26	US: MA	5		0	strawberries, blueberries	(50)	
2006	O026	Japan	16	0	0	Food, restaurant	(185)	served at one restaurant
2006	O026:H11	Japan	33	0	0	Person-person	(247)	nursery school
2006	O080:H- & O145	Germany	59	1		Milk, unpasteurized	(65)	camp
2006	O103:H2	Japan	12	0	0	Person-person	(192)	nursery school
2006	O103:H25	Norway	18	10	1	Mutton, sausage	(106;233;2 36)	
2006	O121:H19	US: UT	69	4		Lettuce	(176)	Wendy's restaurant
2007	O026 & O145	Belgium	13	5	0	ice cream	(60)	contamination by food handlers ?
2007	O026	US: IA	3			Water, pool	(176)	
2007	O026:H-	Japan	29			Person-person	(7)	nursery school
2007	O026:H11	Denmark	20	0	0	Sausage	(74;75)	
2007	O026:H11	Japan	19			Person-person	(7)	nursery school
2007	O026:H11	Japan	19			Person-person	(7)	nursery school
2007	O026:H11	Japan	31			Person-person	(7)	nursery school
2007	O055:H80	Australia	3	1		Animal contact	(110)	kangaroos and koalas at an animal sanctuary; also infected with <i>S. Virchow</i>
2007	O103:H11	Japan	20			Unknown	(7)	old folks home
2007	O111:H-	Japan	8			Person-person	(7)	kindergarten
2007	O111:H-	Japan	28			Unknown	(7)	nursery school
2007	O111:H-	Japan	22			Person-person	(7)	nursery school
2007	O111	US: ME	8			Food	(203)	
2007	O111	US: ND	6			Person-person	(203)	
2007	O111:NM	US: ND	23	0	0	Beef, ground	(195)	wedding reception
2008	O111:NM	US: OK	341	25	1	Food, restaurant	(203)	traced to 1 restaurant

2008	O026:H11	Japan	91			Food, high school	(8)	high school
2008	O026:H11	Japan	27			Person-person	1(8)	nursery school
2008	O026:H11	Japan	52			Unknown	(8)	school excursion
2008	O026:H11	Japan	8			Person-person	(8)	nursery school
2008	O026:H11	Japan	84			Person-person	(8)	kindergarten
2008	O026:H11	Japan	10			Person-person	(8)	nursery school
2008	O026:H11	Japan	11			Person-person	(8)	nursery school
2008	O111:H-	Japan	67			Unknown	(8)	hospital
2008	O111:H-	Japan	13			Unknown	(8)	
2008	O026:H-	Japan	32			Person-person	(8)	nursery school
2008	O026:H-	Japan	14			Person-person	(8)	nursery school
2008	O145:H-	Japan	13			Person-person	(8)	nursery school
2008	O111:H-	Japan	61			Person-person	(8)	nursery school
2008	O115:H NM	Japan	2			Chicken or egg	(76)	
2008	Unknown not O157	Australia	7			Water drinking	(210)	camp
2009	O121	Japan	31			Person-person	(8)	nursery school

Figure 1. Vehicles Associated with Outbreaks of Non-O157:H7 STEC





Figure 2. Vehicles Associated with Outbreak Cases of Non-O157:H7 STEC



References

1. Akiba Y, Kimura T, Takagi M, Akimoto T, Mitsui Y, Ogasawara Y, and Omichi M. 2005. Outbreak of enterohemorrhagic *Escherichia coli* O121 among school children exposed to cattle in a ranch for public education on dairy farming. *Jap J Infect Dis* 58:190-192.
2. Allerberger F, Friedrich AW, Grif K, Dierich MP, Dornbusch HJ, Mache CJ, Nachbaur E, Freilinger M, Rieck P, Wagner M, Caprioli A, Karch H, and Zimmerhack LB. 2003. Hemolytic-uremic syndrome associated with enterohemorrhagic *Escherichia coli* O26:H infection and consumption of unpasteurized cow's milk. *Int J Infect Dis* 7:42-45.
3. Alpers K, Werber D, Frank C, Koch J, Friederich AW, Karch H, An Der Heiden M, Prager R, Fruth A, Bielaszewska M, Morlock G, Heissenhuber A, Diedler A, Gerber A, and Ammon A. 2009. Sorbitol-fermenting enterohaemorrhagic *Escherichia coli* O157:H- causes another outbreak of haemolytic uraemic syndrome in children. *Epidemiol Infect* 137:389-395.
4. Ammon A. 1997. Surveillance of enterohaemorrhagic *E. coli* (EHEC) infections and haemolytic uraemic syndrome (HUS) in Europe. *Euro Surveill* 2:p113.
5. Ammon A, Petersen LR, and Karch H. 1999. A large outbreak of hemolytic uremic syndrome caused by an unusual sorbitol-fermenting strain of *Escherichia coli* O157:H-. *J Infect Dis* 179:1274-1277.
6. Anderson RC, Carr MA, Miller RK, King DA, Carstens GE, Genovese KJ, Callaway TR, Edrington TS, Jung YS, McReynolds JL. 2005. Effects of experimental chlorate preparations as feed and water supplements on *Escherichia coli* colonization and contamination of beef cattle and carcasses. *Food Microbiol* 22:439-447.
7. Anon. 2008. Enterohemorrhagic *Escherichia coli* infection in Japan as of April 2008. *Infect Agents Surveill Rep* 29:117-118.
8. Anon. 2009. Enterohemorrhagic *Escherichia coli* infection in Japan as of April 2009. *Infect Agents Surveill Rep* 30:119-120.
9. Auvray F, Lecureuil C, Dilasser F, Taché J, and Derzelle S. 2009. Development of a real-time PCR assay with an internal amplification control for the screening of shiga toxin-producing *Escherichia coli* in foods. *Lett Appl Microbiol* 48:554-559.
10. Aymerich T, Picouet PA, and Monfort JM. 2008. Decontamination technologies for meat products. *Meat Sci* 78:114-129.
11. Bach SJ, Mcallister TA, Veira DM, Gannon VPJ, and Holley RA. 2003. Effect of bacteriophage DC22 on *Escherichia coli* O157:H7 in an artificial rumen system (Rusitec) and inoculated sheep. *Anim Res* 52:89-101.
12. Banatvala N, Debeukelaer MM, Griffin PM, Barrett TJ, Greene KD, Green JH, and Wells JG. 1996. Shiga-like toxin-producing *Escherichia coli* O111 and associated hemolytic-uremic syndrome - a family outbreak. *Pediatr Infect Dis J* 15:1008-1011.
13. Barkocy-Gallagher GA, Arthur TM, Rivera-Betancourt M, Nou XW, Shackelford SD, Wheeler TL, and Koochmarai M. 2003. Seasonal prevalence of shiga toxin-producing *Escherichia coli*, including O157:H7 and non-O157 serotypes, and *Salmonella* in commercial beef processing plants. *J Food Prot* 66:1978-1986.
14. Barlow RS, Gobius KS, and Desmarchelier PM. 2006. Shiga toxin-producing *Escherichia coli* in ground beef and lamb cuts: results of a one-year study. *Int J Food Microbiol* 111:1-5.
15. Baylis CL. 2008. Growth of pure cultures of verocytotoxin-producing *Escherichia coli* in a range of enrichment media. *J Appl Microbiol* 105:1259-1265.

16. Baylis CI. 2009. Raw milk and raw milk cheeses as vehicles for infection by verocytotoxin-producing *Escherichia coli*. *Int J Dairy Technol* 62(3):293-307.
17. Baylis CL, MacPhee S, Robinson A J, Griffiths R, Lilley K, and Betts RP. 2004. Survival of *Escherichia coli* O157:H7, O111:H- and O26:H11 in artificially contaminated chocolate and confectionery products. *Int J Food Microbiol* 96:35-48.
18. Bergholz TM and Whittam TS. 2007. Variation in acid resistance among enterohaemorrhagic *Escherichia coli* in a simulated gastric environment. *J Appl Microbiol* 102:352-362.
19. Bergmire-Sweat D, Marengo L, Pendergrass P, Hendricks K, Garcia M, Drumgoole R, Baldwin T, Kingsley K, Walsh B, Lang S, Prine L, Busby T, Trujillo L, Perrotta D, Hathaway A, Jones B, and Jaiyeola A. 2000. *Escherichia coli* O111:H8 outbreak among teenage campers - Texas, 1999 . *Morbidity and Mortality Weekly Report* 49:321-324.
20. Berry ED , Barkocy-Gallagher GA, Siragusa GR. 2004. Stationary-phase acid resistance and injury of recent bovine *Escherichia coli* O157 and non-O157 biotype I *Escherichia coli* isolates. *J Food Prot* 67:583-590.
21. Bettelheim KA. 2007. The non-O157 shiga-toxigenic (verocytotoxigenic) *Escherichia coli*; under-rated pathogens. *Crit Rev Microbiol* 33:67-87.
22. Beutin L, Geier D, Zimmermann S, and Karch H. 1995. Virulence markers of shiga-like toxin-producing *Escherichia coli* strains from healthy domestic animals of different species. *J Clin Microbiol* 33:631-635.
23. Beutin L, Jahn S, and Fach P. 2009. Evaluation of the 'GeneDisc' real-time PCR system for detection of enterohaemorrhagic *Escherichia coli* (EHEC) O26, O103, O111, O145 and O157 strains according to their virulence markers and their O- and H-antigen-associated genes. *J Applied Microbiol* 106 :1122-1132.
24. Beutin L, Steinrück, Krause G, Steege K, Haby S, Hultsch G, and Appel B. 2007. Comparative evaluation of the Ridascreen® verotoxin enzyme immunoassay for detection of shiga-toxin producing strains of *Escherichia coli* (STEC) from food and other sources. *J Appl Microbiol* 102:630-639.
25. Beutin L and Strauch E. 2007. Identification of sequence diversity in the *Escherichia Coli* flic genes encoding flagellar types H8 and H40 and its use in typing of shiga toxin-producing *E. coli* O8, O22, O111, O174, and O179 strains. *J Clin Microbiol* 45:333-339.
26. Bhat MA, Nishikawa Y, and Wani SA. 2008. Prevalence and virulence gene profiles of shiga toxin-producing *Escherichia coli* and enteropathogenic *Escherichia coli* from diarrhoeic and healthy lambs in India. *Small Rumin Res* 75:65-70.
27. Bielaszewska M, Srámková L, Janda J, Bláhová K, and Ambrožová H. 1990. Verotoxigenic (enterohaemorrhagic) *Escherichia coli* in infants and toddlers in Czechoslovakia. *Infection* 18:352-356.
28. Birchard K. 1999. *E. coli* O26 outbreak reported in Ireland. *Lancet* 354:1706.
29. Blanco J , Blanco M, Blanco JE, Mora A, Gonzalez EA, Bernardez MI, Alonso MP, Coira A, Rodriguez A, Rey J, Alonso JA, and Usera MA. 2003. Verotoxin-producing *Escherichia coli* in Spain: Prevalence, serotypes, and virulence genes of O157:H7 and non-O157 VTEC in ruminants, raw beef products, and humans. *Exp Biol Med* 228:345-351.
30. Blanco M, Blanco JE, Mora A, Rey J, Alonso JM, Hermoso M, Hermoso J, Alonso MP, Dahbi G, Gonzalez EA, Bernardez MI, and Blanco J. 2003. Serotypes, virulence genes, and intimin types of Shiga toxin (verotoxin)-producing *Escherichia coli* isolates from healthy sheep in Spain. *J Clin Microbiol* 41:1351-1356.
31. Bosilevac JM, Guerini MN, Brichta-Harhay DM, Arthur TM, and Koohmaraie M. 2007. Microbiological

- characterization of imported and domestic boneless beef trim used for ground beef. *J Food Prot* 70:440-449.
32. Bosilevac JM, Nou X, Barkocy-Gallagher GA, Arthur TM, and Koohmaraie M. 2006. Treatments using hot water instead of lactic acid reduce levels of aerobic bacteria and Enterobacteriaceae and reduce the prevalence of *Escherichia coli* O157:H7 on previsceration beef carcasses. *J Food Prot* 69:1808-13.
 33. Boudailliez B, Berquin P, Mariani-Kurkdjian P, Ilef DD, Cuvelier B, Capek I, Tribout B, Bingen E, and Piussan C. 1997. Possible person-to-person transmission of *Escherichia coli* O111 - associated hemolytic uremic syndrome. *Pediatr Nephrol* 11:36-39.
 34. Brooks JT, Bergmire-Sweet D, Kennedy M, Hendricks K, Garcia M, Marengo L, Wells J, Ying M, Bibb W, Griffin PM, Hoekstra RM, and Friedman CR. 2004. Outbreak of Shiga toxin-producing *Escherichia coli* O111:H8 infections among attendees of a high school cheerleading camp. *Clin Infect Dis* 38:190-198.
 35. Brooks JT, Sowers EG, Wells JG, Greene KD, Griffin PM, Hoekstra RM, and Strockbine NA. 2005. Non-O157 shiga toxin-producing *Escherichia coli* infections in the United States, 1983-2002. *J Infect Dis* 192:1422-1429.
 36. Busch U, Hörmansdorfer S, Schraner S, Huber I, Bogner KH, and Sing A. 2007. Enterohemorrhagic *Escherichia coli* excretion by child and her cat. *Emerg Infect Dis* 13:348-349.
 37. Callaway TR, Carr MA, Edrington TS, Anderson RC, and Nisbet DJ. 2009. Diet, *Escherichia Coli* O157:H7, and cattle: a review After 10 Years. *Current Issues Molecular Biol* 11:67-79.
 38. Cameron AS, Beers MY, Walker CC, Rose N, Aneer E, Manatakis Z, Kirke K, Calder I, Jenkins F, Goldwater PN, Paton A, Paton J, Jureidini K, Hoffman A, Henning P, Hansman D, Lawrence A, Miller R, Ratcliff R, Doyle R, Murray C, Davos D, Cameron P, Seymour-Murray J, Lim I, Lanser J, Selvey L, and Beaton S. 1995. Community outbreak of hemolytic uremic syndrome attributable to *Escherichia coli* O111:NM - South Australia, 1995. *Morbidity Mortality Weekly Rep* 44:550-551, 557, 558.
 39. Capek I and Ilef D. 1993. An unusual epidemic in Oise [Fr]. *Bull Epidemiol Hebdomadaire* 48:221-222.
 40. Caprioli A, Luzzi I, Rosmini F, Resti C, Edefonti A, Perfumo F, Farina C, Goglio A, Gianviti A, and Rizzoni G. 1994. Communitywide outbreak of hemolytic-uremic syndrome associated with non-O157 verocytotoxin-producing *Escherichia coli*. *J Infect Dis* 169:208-211.
 41. Caprioli A, Morabito S, Brugere H, and Oswald E. 2005. Enterohaemorrhagic *Escherichia coli*: emerging issues on virulence and modes of transmission. *Vet Res* 36:289-311.
 42. Carroll KC, Adamson K, Korgenski K, Croft A, Hankemeier R, Daly J, and Park CH. 2003. Comparison of a commercial Reversed Passive Latex Agglutination Assay to an enzyme immunoassay for the detection of shiga toxin-producing *Escherichia coli*. *Eur J Clin Microbiol Infect Dis* 22:689-692.
 43. Centers for Disease Control and Prevention. Bacterial foodborne and diarrheal disease national case surveillance. 2003. <http://www.cdc.gov/foodborneoutbreaks/documents/fbsurvsumm2003.pdf>
 44. Centers for Disease Control and Prevention. Bacterial foodborne and diarrheal disease national case surveillance. 2004. <http://www.cdc.gov/foodborneoutbreaks/documents/fbsurvsumm2004.pdf>
 45. Centers for Disease Control and Prevention. Bacterial foodborne and diarrheal disease national case surveillance. 2005. <http://www.cdc.gov/foodborneoutbreaks/documents/fbsurvsumm2005.pdf>
 46. Centers for Disease Control and Prevention. 2006. Importance of culture confirmation of shiga toxin-producing *Escherichia coli* infection as illustrated by outbreaks of gastroenteritis--New York and North Carolina, 2005. *Morbidity Mortality Weekly Rep* 55(38).

47. Centers for Disease Control and Prevention. 2007. Summary of notifiable diseases - United States, 2005. *Morbidity and Mortality Weekly Report* 54:76.
48. Centers for Disease Control and Prevention. 2008. FoodNet Surveillance Report for 2005. http://www.cdc.gov/foodnet/annual/2005/2005_AR_Report.pdf
49. Centers for Disease Control and Prevention. 2009. Compendium of measures to prevent disease associated with animals in public settings. *Morbidity and Mortality Weekly Report* 58:1-21.
50. Centers for Disease Control and Prevention. Outbreak surveillance data. 2009. http://www.cdc.gov/foodborneoutbreaks/outbreak_data.htm
51. Cho S, Diez-Gonzalez F, Fossler CP, Wells SJ, Hedberg CW, Kaneene JB, Ruegg PL, Warnick LD, and Bender JB. 2006. Prevalence of shiga toxin-encoding bacteria and shiga toxin-producing *Escherichia coli* isolates from dairy farms and county fairs. *Veterinary Microbiology* 118:289-298.
52. Cho S, Fossler CP, Diez-Gonzalez F, Wells SJ, Hedberg CW, Kaneene JB, Ruegg PL, Warnick LD, and Bender JB. 2009. Cattle-level risk factors associated with fecal shedding of shiga toxin-encoding bacteria on dairy farms, Minnesota, USA. *Canadian Journal of Veterinary Research* 73:151-156.
53. Cobbold RN, Davis MA, Rice DH, Szymanski M, Tarr PI, Besser TE, and Hancock DD. 2008. Associations between bovine, human, and raw milk and beef isolates of non-O157 shiga toxin-producing *Escherichia coli* within a restricted geographic area of the United States. *Journal of Food Protection* 71:1023-1027.
54. Combs BG, Wise RP, Tribe IG, Mwanri L, and Raupach JCA. 2003. Investigation of two clusters of shiga toxin-producing *Escherichia coli* cases in South Australia. *Communicable Disease Intelligence* 27:517-519.
55. Cookson AL, Taylor SCS, Bennett J, Thomson-Carter F, and Attwood GT. 2006. Serotypes and analysis of distribution of shiga toxin-producing *Escherichia coli* from cattle and sheep in the Lower North Island, New Zealand. *New Zealand Veterinary Journal* 54:78-84.
56. Coombes BK, Wickham ME, Mascarenhas M, Gruenheid S, Finlay BB, and Karmali MA. 2008. Molecular analysis as an aid to assess the public health risk of non-O157 shiga toxin-producing *Escherichia coli* strains. *Applied and Environmental Microbiology* 74:2153-2160.
57. Cutter CN and Rivera-Betancourt M. 2000. Interventions for the reduction of *Salmonella Typhimurium* DT 104 and Non-O157:H7 enterohemorrhagic *Escherichia coli* on beef surfaces. *Journal of Food Protection* 63:1326-1332.
58. De Paula CJS and Marin JM. 2008. Occurrence of non-O157 shiga toxin-producing *Escherichia coli* in dogs with diarrhea. *Ciencia Rural* 38:1682-1686.
59. De Sablet T, Bertin Y, Vareille M, Girardeau JP, Garrivier A, Gobert AP, and Martin C. 2008. Differential expression of Stx(2) variants in shiga toxin-producing *Escherichia coli* belonging to seropathotypes A and C. *Microbiology* 154:176-186.
60. De Schrijver K, Buvens G, Possé B, Van Den Branden D, Oosterlynck C, De Zutter L, Eilers K, Piérard D, Dierick K, Van Damme-Lombaerts R, Lauwers C, and Jacobs R. 2008. Outbreak of verocytotoxin-producing *E. coli* O145 and O26 infections associated with the consumption of ice cream produced at a farm, Belgium, 2007. *Euro Surveill* 13.
61. Deschênes G, Casenave C, Grimont F, Desenclos JC, Benoit S, Collin M, Baron S, Mariani P, Grimont P, and Nivet H. 1996. Cluster of cases of haemolytic uraemic syndrome due to unpasteurised cheese. *Pediatric Nephrology* 10:203-205.
62. Desrosiers A, Fairbrother JM, Johnson RP, Desautels C, Letellier A, and Quessy S. 2001. Phenotypic and

- genotypic characterization of *Escherichia coli* verotoxin-producing isolates From humans and pigs. J Food Prot 64:1904-1911.
63. Djordjevic SP, Ramachandran V, Bettelheim KA, Vanselow BA, Holst P, Bailey G, and Hornitzky MA. 2004. Serotypes and virulence gene profiles of shiga toxin-producing *Escherichia coli* strains Isolated from feces of pasture-fed and lot-fed sheep. Appl Environ Microbiol 70:3910-3917.
 64. Doyle ME, Archer J, Kaspar CW, and Weiss R. Human illness caused by *E. coli* O157:H7 from food and non-food sources. 2006. http://fri.wisc.edu/briefs/FRIBrief_EcoliO157H7humanillness.pdf
 65. Dreesman J, Pulz M, Röttgers HR, and Mellman A. 2008. Outbreak of EHEC infection associated with raw milk consumption at a holiday camp [Ger]. Epidemiol Bull Robert Koch Inst 16-18.
 66. Duffy G, Walsh C, Blair IS, and McDowell DA. 2006. Survival of antibiotic resistant and antibiotic sensitive strains of *E. coli* O157 and *E. coli* O26 in food matrices. Int J Food Microbiol 109:179-186.
 67. Duris JW, Haack SK, and Fogarty LR. 2009. Gene and antigen markers of shiga-toxin producing *E. coli* from Michigan and Indiana river water: occurrence and relation to recreational water quality criteria. J Environ Qual 38:1878-86.
 68. Durso LM, Bono JL, and Keen JE. 2005. Molecular serotyping of *Escherichia coli* O26:H11. Appl Environ Microbiol 71:4941-4944.
 69. Durso LM, Bono JL, and Keen JE. 2007. Molecular serotyping of *Escherichia coli* O111:H8. J Microbiol Meth 69:381-383.
 70. Döpfer D, Geue L, de Bree J, and de Jong MCM. 2006. Dynamics of verotoxin-producing *Escherichia coli* isolated from German beef cattle between birth and slaughter. Prev Vet Med 73:229-240.
 71. Elliott EJ, Robins-Browne RM, O'Loughlin EV, Bennett-Wood V, Bourke J, Henning P, Hogg GG, Knight J, Powell H, and Redmond D. 2001. Nationwide study of haemolytic uraemic syndrome: clinical, microbiological, and epidemiological features. Arch Dis Child 85:125-131.
 72. Espié E, Grimont F, Mariani-Kurkdjian P, Bouvet P, Haeghebaert S, Filliol I, Loirat C, Decludt B., Minh NNT, Vaillant V, and De Valk H. 2008. Surveillance of hemolytic uremic syndrome in children less than 15 years of age, a system to monitor O157 and non-O157 shiga toxin-producing *Escherichia coli* infections in France, 1996-2006. Pediatr Infect Dis J 27:595-601.
 73. Espié E, Grimont F, Vaillant V, Montet MP, Carle I, Bavai C, De Valk H, and Vernozy-Rozand C. 2006. O148 shiga toxin-producing *Escherichia coli* outbreak: microbiological investigation as a useful complement to epidemiological investigation. Clin Microbiol Infect 12:992-998.
 74. Ethelberg S, Smith B, Torpdahl M, Lisby M, Boel J, Jensen T, and Mølbak K. 2007. An outbreak of verocytotoxin-producing *Escherichia coli* O26:H11 caused by beef sausage, Denmark 2007. Euro Surveill 12.
 75. Ethelberg S, Smith B, Torpdahl M, Lisby M, Boel J, Jensen T, Nielsen EM, and Mølbak K. 2009. Outbreak of non-O157 shiga toxin-producing *Escherichia coli* infection from consumption of beef sausage. Clin Infect Dis 48:E78-E81.
 76. Etoh Y, Murakami K, Ichihara S, Sera N, Hamasaki M, Takenaka S, Horikawa K, Kawano K, Takeishi T, Kuwana Y, Inoue A, Nagatsu Y, Hira Y, Takahashi M, and Ito K. 2009. Isolation of shiga toxin 2f-producing *Escherichia coli* (O115:HNM) from an adult symptomatic patient in Fukuoka Prefecture, Japan. Jap J Infect Dis 62:315-317.
 77. Evans J, Knight HI, Smith AW, Pearce MC, Hall M, Foster G, Low JC, and Gunn GJ. 2008. Cefixime-

- tellurite rhamnose MacConkey agar for isolation of verocytotoxin-producing *Escherichia coli* serogroup O26 from Scottish cattle and sheep faeces. *Lett Appl Microbiol* 47:148-152.
78. Farooq S, Hussain I, Mir MA, Bhat MA, and Wani SA. 2009. Isolation of atypical enteropathogenic *Escherichia Coli* and shiga toxin 1 and 2f-producing *Escherichia Coli* from avian species in India. *Lett Appl Microbiol* 48:692-697.
 79. Feng P, Weagant SD, and Monday SR. 2001. Genetic analysis for virulence factors in *Escherichia coli* O104:H21 that was implicated in an outbreak of hemorrhagic colitis. *J Clin Microbiol* 39:24-28.
 80. Fischer JR, Zhao T, Doyle MP, Goldberg MR, Brown CA, Sewell CT, Kavanaugh DM, and Bauman CD. 2001. Experimental and field studies of *Escherichia coli* O157:H7 in white-tailed deer. *Appl Environ Microbiol* 67:1218-1224.
 81. Frank C, Kapfhammer S, Werber D, Stark K, and Held L. 2008. Cattle density and shiga toxin-producing *Escherichia coli* infection in Germany: increased risk for most but not all serogroups. *Vector-borne Zoonotic Dis* 8:635-643.
 82. Fratamico PM, Bagi LK, Bush EJ, and Solow BT. 2004. Prevalence and characterization of shiga toxin-producing *Escherichia coli* in swine feces recovered in the National Animal Health Monitoring System's Swine 2000 study. *Appl Environ Microbiol* 70:7173-7178.
 83. Fratamico PM, Debroy C, Liu YH. 2009. The DNA sequence of the *Escherichia coli* O22 O-antigen gene cluster and detection of pathogenic strains belonging to *E. coli* serogroups O22 and O91 by multiplex PCR assays targeting virulence genes and genes in the respective O-antigen gene clusters. *Food Anal Meth* 2:169-179.
 84. Fratamico PM, DebRoy C, Miyamoto T, and Liu YH. 2009. PCR Detection of enterohemorrhagic *Escherichia coli* O145 in food by targeting genes in the *E. coli* O145 O-antigen gene cluster and the shiga toxin 1 and shiga toxin 2 Genes. *Foodborne Path Dis* 6:605-611.
 85. Fremaux B, Delignette-Muller ML, Prigent-Combaret C, Gleizal A, and Vernozy-Rozand C. 2007. Growth and survival of non-O157:H7 shiga-toxin-producing *Escherichia coli* in cow manure. *J Appl Microbiol* 102:89-99.
 86. Fremaux B, Prigent-Combaret C, Delignette-Muller ML, Dothal M, and Vernozy-Rozand C. 2007. Persistence of shiga toxin-producing *Escherichia coli* O26 in cow slurry. *Lett Appl Microbiol* 45:55-61.
 87. Fremaux B, Prigent-Combaret C, Delignette-Muller ML, Mallen B, Dothal M, Gleizal A, and Vernozy-Rozand C. 2008. Persistence of shiga toxin-producing *Escherichia coli* O26 in various manure-amended soil types. *J Appl Microbiol* 104:296-304.
 88. Froehlich J, Baljer G, and Menge C. 2009. Maternally and naturally acquired antibodies to shiga toxins in a cohort of calves shedding shiga-toxigenic *Escherichia coli*. *Appl Environ Microbiol* 75:75 (11) 3695-3704.
 89. Fukushima H, Hoshina K, and Gomyoda M. 1999. Long-term survival of shiga toxin-producing *Escherichia coli* O26, O111, and O157 in cattle feces. *Appl Environ Microbiol* 65:5177-5181.
 90. Förster M, Sievert K, Messler S, Klimpel S, and Feffer K. 2009. Comprehensive study on the occurrence and distribution of pathogenic microorganisms carried by synanthropic flies caught at different rural locations in Germany. *J Med Entomol* 46:1164-1166.
 91. Garcia A and Fox JG . 2003. The rabbit as a new reservoir host of enterohemorrhagic *Escherichia coli*. *Emerg Infect Dis* 9:1592-1597.

92. Gareis M, Pichner R, Brey N, and Steinrueck H. 2000. Shedding of verotoxigenic *E. coli* by healthy staff of a food producing company. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*: 43 (10) 781-787 43:781-787.
93. Garvey P, McKeown P, Carroll A, and McNamara E. 2009. Epidemiology of verotoxigenic *E. coli* in Ireland, 2008. *Epi-Insight* 10.
94. Gavin PJ, Peterson LR, Pasquariello AC, Blackburn J, Hamming MG, Kuo KJ, and Thomson RB. 2004. Evaluation of performance and potential clinical impact of Prospect Shiga Toxin *Escherichia coli* Microplate Assay for detection of shiga toxin-Producing *E. coli* in stool samples. *J Clin Microbiol* 42:1652-1656.
95. Gilbert RA, Denman SE, Padmanabha J, Fegan N, Ajmi D, and McSweeney CS. 2008. Effect of diet on the concentration of complex shiga toxin-producing *Escherichia coli* and EHEC virulence genes in bovine faeces, hide and carcass. *Int J Food Microbiol* 121:208-216.
96. Gilbreath JJ, Shields MS, Smith RL, Farrell LD, Sheridan PP, and Spiegel KM. 2009. Shiga toxins, and the genes encoding them, in fecal samples from native Idaho ungulates. *Appl Environ Microbiol* 75:862-865.
97. Goncalves VP and Morin JM. 2007. Fate of non O157 shiga toxigenic *Escherichia coli* in composted cattle manure. *Arquivo Brasileiro De Medicina Veterinaria E Zootecnia* 59:825-831.
98. Gough JM, Conlan LL, Denman SE, Krause DO, Smith WJM, Williamson MA, and McSweeney CS. 2006. Screening of bacteria from the cattle gastrointestinal tract for inhibitory activity against enterohemorrhagic *Escherichia coli* O157:H7, O111:H-, and O26:H11. *J Food Prot* 69:2843-2850.
99. Gould HL, Bopp C, Strockbine N, Atkinson R, Baselski V, Body B, Carey R, Crandall C, Hurd S, Kaplan RAY, Neill M, Shea S, Somsel P, Tobin-D'angelo M, Griffin PM, Gerner-Smidt P. 2009. Recommendations for diagnosis of shiga toxin--producing *Escherichia coli* infections by clinical laboratories. *MMWR Recomm Rep* 58:1-14.
100. Gourmelon M, Montet MP, Lozach S, Le Mennec C, Pommepuy M, Beutin L, and Vernozy-Rozand C. 2006. First isolation of Shiga toxin 1d producing variant strains in shellfish from coastal areas in France. *J Appl Microbiol* 100:85-97.
101. Grant MA, Mogler MA, and Harris DL. 2009. Comparison of enrichment procedures for shiga toxin-producing *Escherichia coli* in wastes from commercial swine farms. *J Food Prot* 72:1982-1986.
102. Grossmann K, Weniger B, Baljer G, Brenig B, and Wieler LH. 2005. Racing, ornamental and city pigeons carry Shiga toxin producing *Escherichia coli* (STEC) with different Shiga toxin subtypes, urging further analysis of their epidemiological role in the spread of STEC. *Berlin Munch Tierarzt Wochenschr* 118:456-463.
103. Grys TE, Sloan LM, Rosenblatt JE, and Patel R. 2009. Rapid and sensitive detection of shiga toxin-producing *Escherichia coli* from nonenriched stool specimens by real-time PCR in comparison to enzyme immunoassay and culture. *J Clin Microbiol* 47:2008-2012.
104. Grzadkowska D and Griffiths MW. 2001. Cryotolerance of *Escherichia coli* O157:H7 in laboratory media and food. *J Food Sci* 66:1169-1173.
105. Gu J, Liu YQ, Yu S, Wang HG, Wang QX, Yi Y, Zhu FC, Yu XJ, Zou QM, and Mao XH. 2009. Enterohemorrhagic *Escherichia coli* trivalent recombinant vaccine containing EspA, intimin and Stx2 Induces strong humoral immune response and confers protection in mice. *Microbes Infect* 11:835-841.
106. Gudmundsdottir H, Krogvold L, Henrichsen T, Draganov B, Bangstad HJ, Bjerre A, and Os I. 2007. Dialysis in hemolytic uremic syndrome in small children. report of a major outbreak of shiga-toxin-producing strain of

- E. coli* O103:H25 in Norway. *Nephrol Dialysis Transplant* 22 Suppl. 6:51.
107. Guion, C. E., T. J. Ochoa, C. M. Walker, F. Barletta, and T. G. Cleary. 2008. Detection of Diarrheagenic *Escherichia Coli* by Use of Melting-Curve Analysis and Real-Time Multiplex PCR. *J Clin Microbiol* 46:1752-1757.
 108. Gyles CL. 2007. Shiga toxin-producing *Escherichia coli*: an overview. *J Anim Sci* 85:E45-E62.
 109. Hall LM, Evans J, Smith AW, Pearce MC, Knight HI, Foster G, Low JC, and Gunn GJ. 2006. Sensitivity of an immunomagnetic-separation-based test for detecting *Escherichia coli* O26 in bovine feces. *Appl Environ Microbiol* 72:7260-7263.
 110. Hanna JN, Humphreys JL, Ashton SE, and Murphy DM. 2007. Haemolytic uremic syndrome associated with a family cluster of enterohaemorrhagic *Escherichia coli*. *Commun Dis Intell* 31:300-303.
 111. Hara-Kudo Y, Konishi N, Ohtsuka K, Hirramatsu R, Tanaka H, Konum H, and Takatori K. 2008. Detection of verotoxigenic *Escherichia coli* O157 and O26 in food by plating methods and LAMP method: a collaborative study. *Int J Food Microbiol* 122:156-161.
 112. Hara-Kudo Y, Nemoto J, Ohtsuka K, Segawa Y, Takatori K, Kojima T, and Ikedo M. 2007. Sensitive and Rapid detection of vero toxin-producing *Escherichia coli* using loop-mediated isothermal mmplication. *J Med Microbiol* 56:398-406.
 113. Hara-Kudo Y, Niizuma J, Goto I, Iizuka S, Kaji Y, Kamakura K, Suzuki S, and Takatori K. 2008. Surveillance of shiga toxin-producing *Escherichia coli* in beef with effective procedures, independent of serotype. *Foodborne Path Dis* 5:97-103.
 114. Hashimoto H, Mizukoshi K, Nishi M, Kawakita T, Hasui S, Kato Y, Ueno Y, Takeya R, Okuda N, and Takeda T. 1999. Epidemic of gastrointestinal tract infection including hemorrhagic colitis attributable to shiga toxin 1-producing *Escherichia coli* O118:H2 at a junior high school in Japan. *Pediatrics* 103.
 115. Haus-Cheymol R, Espie E, Che D, Vaillant V, DeValk H, and Desenclos JC. 2006. Association between indicators of cattle density and incidence of paediatric haemolytic-uraemic syndrome (HUS) in children under 15 years of age in France between 1996 and 2001: an ecological study. *Epidemiol Infect* 134:712-718.
 116. Hazarika RA, Singh DK, Kapoor KN, Agarwal RK, Pandey AB, and Rajkumar DN. 2004. Detection and characterization of verotoxin-producing *Escherichia coli* (VTEC) isolated from buffalo meat. *J Food Safety* 24:281-290.
 117. Hazarika RA, Singh DK, Kapoor KN, K. Bhilegaonkar, Agarwal RK, and Malik SVS. 2003. Survivability of vero toxic *Escherichia coli* O111 against sodium benzoate and potassium sorbate added to simulating media for beef gravy. *J Vet Public Health* 1:103-112.
 118. Heinikainen S, Pohjanvirta T, Eklund M, Siitonen A, and Pelkonen S. 2007. Tracing shigatoxigenic *Escherichia coli* O103, O145, and O174 infections from farm residents to cattle. *J Clin Microbiol* 45:3817-3820.
 119. Henderson H. 2008. Direct and indirect zoonotic transmission of shiga toxin-producing *Escherichia coli*. *J Am Vet Med Assoc* 232:848-859.
 120. Heuvelink AE, Zwartkruis-Nahuis JTM, van den Biggelaar FLAM, van Leeuwen WJ, and de Boer E. 1999. Isolation and characterization of verocytotoxin-producing *Escherichia coli* O157 from slaughter pigs and poultry. *Int J Food Microbiol* 52:67-75.
 121. Hiramatsu R, Matsumoto M, Sakae K, and Miyazaki Y. 2005. Ability of Shiga toxin-producing *Eschetichia*

- coli* and *Salmonella* spp. to survive in a desiccation model system and in dry foods. *Appl Environ Microbiol* 71:6657-6663.
122. Hiruta N, Murase T, and Okamura N. 2001. An outbreak of diarrhoea due to multiple antimicrobial-resistant shiga toxin-producing *Escherichia coli* O26:H11 in a nursery. *Epidemiol Infect* 127:221-227.
 123. Hoshina K, Itagaki A, Seki R, Yamamoto K, Masuda S, Muku T, and Okada N. 2001. Enterohemorrhagic *Escherichia coli* O26 outbreak caused by contaminated natural water supplied by facility owned by local community. *Jap J Infect Dis* 54:247-248.
 124. Hu J, Green DA, Swoveland J, Grant M, and Boyle DS. 2009. Preliminary evaluation of a procedure for improved detection of shiga toxin-producing *Escherichia coli* in fecal specimens. *Diagn Microbiol Infect Dis* 65:21-26.
 125. Hughes LA, Bennett M, Coffey P, Elliott J, Jones TR, Jones RC, Lahuerta-Marin A, McNiffe K, Norman D, Williams NJ, and Chantrey J. 2009. Risk factors for the occurrence of *Escherichia coli* virulence genes *eae*, *stx1* and *stx2* in wild bird populations. *Epidemiol Infect* 137:1574-1582.
 126. Hugo AA, Kakisu E, De Antoni GL, and Perez PF. 2008. Lactobacilli antagonize biological effects of enterohaemorrhagic *Escherichia coli* in vitro. *Lett Appl Microbiol* 46:613-619.
 127. Hundy RL and Cameron S. 2004. Risk factors for sporadic human infection with shiga toxin-producing *Escherichia coli* in South Australia. *Commun Dis Intelligence* 28:74-79.
 128. Hung VK and Cornick NA. 2008. Prevalence and genetic profiles of shiga toxin-producing *Escherichia coli* strains isolated from buffaloes, cattle, and goats in central Vietnam. *Vet Microbiol* 126:356-363.
 129. Hussein HS. 2007. Prevalence and pathogenicity of shiga toxin-producing *Escherichia coli* in beef cattle and their products. *J Animal Sci* 85 Suppl. 1:E63-E72.
 130. Hussein HS and Bollinger LM. 2005. Prevalence of Shiga toxin-producing *Escherichia coli* in beef. *Meat Sci* 71:676-689.
 131. Hussein HS and Bollinger LM. 2005. Prevalence of Shiga toxin-producing *Escherichia coli* in beef cattle. *J Food Prot* 68:2224-2241.
 132. Hussein HS, Bollinger LM, and Hall MR. 2008. Growth and enrichment medium for detection and isolation of shiga toxin-producing *Escherichia coli* in cattle feces. *J Food Prot* 71:927-933.
 133. Hussein HS and Sakuma T. 2005. Prevalence of Shiga toxin-producing *Escherichia coli* in dairy cattle and their products. *J Dairy Sci* 88:450-465.
 134. Hussein HS and Sakuma T. 2005. Shiga toxin-producing *Escherichia coli*: pre- and postharvest control measures to ensure safety of dairy cattle products. *J Food Prot* 68:199-207.
 135. Hussein HS, Thran BH, and Glimp HA. 2003. Verotoxin-producing *Escherichia coli* in sheep grazing an irrigated pasture or arid rangeland forages. *Exp Biol Med* 228:358-364.
 136. Iizuka S, Tsunomori Y, Tabara K, Tsuda K, and Fukuma T. 2005. An outbreak of mixed infection of enterohemorrhagic *Escherichia coli* O26:H11 and norovirus genogroup II at a kindergarten in Shimane, Japan. *Jap J Infect Dis* 58:329-330.
 137. Imamovic L, Jofre J, Schmidt H, Serra-Moreno R, and Muniesa M. 2009. Phage-mediated shiga toxin 2 gene transfer in food and water. *Appl Environ Microbiol* 75:1764-1768.

138. Islam MA, Mondol AS, De Boer E, Beumer RR, Zwietering MH, Talukder KA, and Heuvelink AE. 2008. Prevalence and genetic characterization of shiga toxin-producing *Escherichia coli* isolates from slaughtered animals in Bangladesh. *Appl Environ Microbiol* 74:5414-5421.
139. Jenkins C, Evans J, Chart H, Willshaw GA, and Frankel G. 2008. *Escherichia coli* serogroup O26 - a new look at an old adversary. *J Appl Microbiol* 104:14-25.
140. Jeon BW/Jeong JM, Won GY, Park H, Eo SK, Kang HY, Hur J, and Lee JH. 2006. Prevalence and characteristics of *Escherichia coli* O26 and O111 from cattle in Korea. *Int J Food Microbiol* 110:123-126.
141. Johnsen G, Wasteson Y, Heir E, Berget OI, and Herikstad H. 2001. *Escherichia coli* O157 H7 in faeces from cattle, sheep and pigs in the southwest part of Norway during 1998 and 1999. *Int J Food Microbiol* 65:193-200.
142. Johnson KE, Thorpe CM, Sears CL. 2006. The emerging clinical importance of non-O157 shiga toxin - producing *Escherichia coli*. *Clin Infect Dis* 43:1587-1595.
143. Johnson RP, Clarke RC, Wilson JB, Read SC, Rahn K, Renwick SA, Sandhu KA, Alves D, Karmali MA, Lior H, McEwen SA, Spika JS, and Gyles CL. 1996. Growing concerns and recent outbreaks involving non-O157:H7 serotypes of verotoxigenic *Escherichia coli*. *J Food Prot* 59:1112-1122.
144. Kalchayanand N, Arthur TM, Bosilevac JM, Brichta-Harhay DM, Guerini MN, Shackelford SD, Wheeler TL, and Koohmaraie M. 2007. Microbiological characterization of lamb carcasses at commercial processing plants in the United States. *J Food Prot* 70:1811-1819.
145. Kanki M, Seto K, Sakata J, Harada T, and Kumeda Y. 2009. Simultaneous enrichment of shiga toxin-producing *Escherichia coli* O157 and O26 and *Salmonella* in food samples using universal preenrichment broth. *J Food Prot* 72:2065-2070.
146. Karch H and Bielaszewska M. 2001. Sorbitol-fermenting Shiga toxin-producing *Escherichia coli* O157:H- strains: epidemiology, phenotypic and molecular characteristics, and microbiological diagnosis. *J Clin Microbiol* 39:2043-2049.
147. Karmali MA, Mascarenhas M, Shen SH, Ziebell K, Johnson S, Reid-Smith R, Isaac-Renton J, Clarks C, Rahn K, and Kaper JB. 2003. Association of genomic O(-)island 122 of *Escherichia coli* EDL 933 with verocytotoxin-producing *Escherichia coli* seropathotypes that are linked to epidemic and/or serious disease. *J Clin Microbiol* 41:4930-4940.
148. Karmali MA, Petric M, Lim C, Fleming PC, Arbus GS, and Lior H. 1985. The association between idiopathic hemolytic uremic syndrome and infection by verotoxin-producing *Escherichia coli*. *J Infect Dis* 151:775-782.
149. Kato K, Shimoura R, Nashimura K, Yoshifuzi K, Shiroshita K, Sakurai N, Kodama H, and Kuramoto S. 2005. Outbreak of enterohemorrhagic *Escherichia coli* O111 among high school participants in excursion to Korea. *Jap J Infect Dis* 58:332-333.
150. Kaufmann M, Zweifel C, Blanco M, Blanco JE, Blanco J, Beutin L, and Stephan R. 2006. *Escherichia coli* O157 and non-O157 Shiga toxin-producing *Escherichia coli* in fecal samples of finished pigs at slaughter in Switzerland. *J Food Prot* 69:260-266.
151. Keen JE, Wittum JE, Dunn JR, Bono JL, and Durso LM. 2006. Shiga-toxigenic *Escherichia coli* O157 in agricultural fair livestock, United States. *Emerg Infect Dis* 12:780-786.
152. Keene WE, Sazie E, Kok J, Rice DH, Hancock DD, Balan VK, Zhao T, and Doyle MP. 1997. An outbreak of *Escherichia coli* O157:H7 infections traced to jerky made from deer meat. *J Am Med Assoc* 277:1229-1231.

153. Kehl KS , Havens P, Behnke CE, and Acheson DW K. 1997. Evaluation of the Premier EHEC assay for detection of shiga toxin-producing *Escherichia coli*. J Clin Microbiol 35:2051-2054.
154. Khan AB , Naim A, Orth D, Grif K, Mohsin M, Prager R, Dierich MP, and Wurzner R. 2009. Serine protease espP subtype alpha, but not beta or gamma, of shiga toxin-producing *Escherichia coli* is associated with highly pathogenic serogroups . Int J Med Microbiol 299:247-254.
155. Kijima-Tanaka M, Ishihara K, Kojima A, Morioka A, Nagata R, Kawanishi M, Nakazawa M, Tamura Y, and Takahashi T. 2005. A national surveillance of Shiga toxin-producing *Escherichia coli* in food-producing animals in Japan. J Vet Med Ser B 52:230-237.
156. Kobayashi H, Kanazaki M, Hata E, and Kubo M. 2009. Prevalence and characteristics of eae- and stx-positive strains of *Escherichia coli* from wild birds in the immediate environment of Tokyo Bay. Appl Environ Microbiol 75:292-295.
157. Kobayashi H, Kanazaki M, Ogawa T, Iyoda S, and Hara-Kudo Y. 2009. Changing prevalence of O-serogroups and antimicrobial susceptibility among STEC strains isolated from healthy dairy cows over a decade in Japan between 1998 and 2007. J Vet Med Sci 71:363-366.
158. Koohmaraie M, Arthur TM, Bosilevac JM, Guerini M, Shackelford SD, and Wheeler TL. 2005. Post-harvest interventions to reduce/eliminate pathogens in beef. Meat Sci 71:79-91.
159. Kuda T, Yano T, and Kuda MT. 2008. Resistances to benzalkonium chloride of bacteria dried with food elements on stainless steel surface. Lwt-Food Sci Technol 41:988-993.
160. Kumar HS, Otta SK, Karunasagar I, and Karunasagar I. 2001. Detection of shiga-toxigenic *Escherichia coli* (STEC) in fresh seafood and meat marketed in Mangalore, India by PCR. Lett Appl Microbiol 33:334-338.
161. Lainhart W, Stolfa G, and Koudelka GB. 2009. Shiga toxin as a bacterial defense against a eukaryotic predator, *Tetrahymena thermophila*. J Bacteriol 191:5116-5122.
162. Large TM, Walk ST, and Whittam TS. 2005. Variation in acid resistance among shiga toxin-producing clones of pathogenic *Escherichia coli*. Appl Environ Microbiol 71:2493-2500.
163. Lathrop S, Edge K, and Baretta J. 2009. Shiga-toxin producing *Escherichia coli*, New Mexico, USA, 2004-2007. Emerg Infect Dis 15:1289-1291.
164. Leclercq A and Mahillon J . 2003. Farmed rabbits and ducks as vectors for VTEC O157:H7. Vet Rec 152:723-4.
165. Lee SH, Levy DA, Craun GF, Beach MJ, and Calderon RL. 2002. Surveillance for waterborne disease outbreaks - United States - 1999-2000. Morbid Mortal Weekly Rep 51:1-28.
166. Leitch ECM and Stewart CS. 2002. Susceptibility of *Escherichia coli* O157 and non-O157 isolates to lactate. Lett Appl Microbiol 35:176-180.
167. LeJeune JT and Davis MA. 2004. Outbreaks of zoonotic enteric disease associated with animal exhibits. J Am Vet Med Assoc 224:1440-1445.
168. Leotta GA, Deza N, Origlia J, Toma C, Chinen I, Miliwebsky E, Iyoda S, Sosa-Estani S, and Rivas M. 2006. Detection and characterization of shiga toxin-producing *Escherichia coli* in captive non-domestic mammals. Vet Microbiol 118:151-157.
169. Liu J, Sun Y, Feng SZ, Zhu LW, Guo XJ, and Qi C. 2009. Towards an attenuated enterohemorrhagic *Escherichia coli* O157:H7 vaccine characterized by a deleted *ler* gene and containing a pathogenic shiga

* Corresponding author: M. Ellin Doyle, Ph.D., medoyle@wisc.edu
 Food Research Institute, UW–Madison, www.fri.wisc.edu/

- toxins. *Vaccine* 27:5929-5935.
170. Liu WC, Shaw DJ/Matthews L, Hoyle DV, Pearce MC, Yates CM, Low JC, Amyes SGB, Gunn GJ, and Woolhouse MEJ. 2007. Modelling the epidemiology and transmission of verocytotoxin-producing *Escherichia coli* serogroups O26 and O103 in two different calf cohorts. *Epidemiol Infect* 135:1316-1323.
 171. Lockary VM, Hudson RF, and Ball CL. 2007. Shiga toxin-producing *Escherichia coli*, Idaho. *Emerg Infect Dis* 13:1262-1264.
 172. Lorusso V, Dambrosio A, Normanno G, Quaglia NC, Celano GV, Germinario GL, Lucifora G, and Salandra GLA. 2009. Duplex PCR for the identification and characterization of *Escherichia coli* O26 VTEC. *Industrie Alimentari* 48:44-48.
 173. Lorusso V, Dambrosio A, Quaglia NC, Parisi A, La Salandra G, Lucifora G, Mula G, Virgilio S, Carosielli L, Rella A, Dario M, and Normanno G. 2009. Verocytotoxin-producing *Escherichia coli* O26 in raw water buffalo (*Bubalus bubalis*) milk products in Italy. *J Food Prot* 72:1705-1708.
 174. Madic J, Lecureuil C, Dilasser F, Derzelle S, Jamet E, Fach P, and Auvray F. 2009. Screening of food raw materials for the presence of shiga toxin-producing *Escherichia coli* O91:H21. *Lett Appl Microbiol* 48:447-451.
 175. Manna SK, Das R, and Manna C. 2008. Microbiological quality of finfish and shellfish with special reference to shiga toxin-producing *Escherichia coli* O157. *J Food Sci* 73:M283-M286.
 176. Marler Clark. *E. coli* blog. 2007. <http://www.ecoliblog.com/articles/e-coli-outbreaks/>
 177. McCarthy TA, Barrett NL, Hadler JL, Salsbury B, Howard RT, Dingman DW, Brinkman CD, Bibb WF, and Cartter ML. 2001. Hemolytic-uremic syndrome and *Escherichia coli* O121 at a Lake in Connecticut, 1999. *Pediatrics* 108:e59.
 178. McPherson M, Kirk M, Raupach J, and Koehler A. 2008. Risk factors of sporadic human infection of shiga toxin producing *Escherichia coli* O157 and non-O157 in Australia. *Int J Infect Dis* 12:E217.
 179. Mellmann A, Bielaszewska M, and Karch H. 2009. Intrahost genome alterations in enterohemorrhagic *Escherichia coli*. *Gastroenterology* 136:1925-1938.
 180. Mellmann A, Fruth A, Friedrich AW, Wieler LH, Harmsen D, Werber D, Middendorf B, Bielaszewska M, and Karch H. 2009. Phylogeny and disease association of shiga toxin-producing *Escherichia coli* O91. *Emerg Infect Dis* 15:1474-1477.
 181. Mercado EC, Rodriguez SM, Elizondo AM, Marcoppido G, and Parreno V. 2004. Isolation of shiga toxin-producing *Escherichia coli* from a South American camelid (*Lama guanicoe*) with diarrhea. *J Clin Microbiol* 42:4809-4811.
 182. Miko A, Pries K, Haby S, Steege K, Albrecht N, Krause G, and Beutin L. 2009. Assessment of shiga toxin-producing *Escherichia coli* isolates from wildlife meat as potential pathogens for humans. *Appl Environ Microbiol* 75:6462-6470.
 183. Minnesota Dept. of Health. *Escherichia coli* O157 infection and hemolytic uremic syndrome (HUS), 2001. 2002. <http://www.health.state.mn.us/divs/idepc/newsletters/dcn/sum01/ecoli.html>
 184. Misselwitz J, Karch H, Bielaszewska M, John U, Ringelmann F, Rönnefarth G, and Patzer L. 2003. Cluster of hemolytic-uremic syndrome caused by shiga toxin-producing *Escherichia coli* O26:H11. *Pediatr Infect Dis J* 22:349-354.

185. Miyajima Y, Takahashi M, Eguchi H, Honma M, Tanahashi S, Matui Y, Kobayashi G, Tanaka M, Higuchi T, and Takeuchi Y. 2007. Outbreak of enterohemorrhagic *Escherichia coli* O26 in Niigata City, Japan. *Jap J Infect Dis* 60:238-239.
186. Molina PM, Sanz ME, Lucchesi PMA, Padola NL, and Parma AE. 2005. Effects of acidic broth and juices on the growth and survival of verotoxin-producing *Escherichia coli* (VTEC). *Food Microbiol* 22:469-473.
187. Monday SR, Beisaw A, and Feng PCH. 2007. Identification of shiga toxin-producing *Escherichia coli* seropathotypes A and B by multiplex PCR. *Molec Cell Probes* 21:308-311.
188. Montet MP, Christeans S, Thevenot D, Coppet V, Ganet S, Muller MLD, Duniere L, Miszczycha S, and Vernozy-Rozand C. 2009. Fate of acid-resistant and non-acid resistant shiga toxin-producing *Escherichia coli* strains in experimentally contaminated French fermented raw meat sausages. *Int J Food Microbiol* 129:264-270.
189. Moore K, Damrow T, Abbott DO, and Jankowski S. 1995. Outbreak of acute gastroenteritis attributable to *Escherichia coli* serotype O104:H21--Helena, Montana, 1994. *Morbidity and Mortality Weekly Report* 44:501-3.
190. Morabito S, Dell'Omo G, Agrimi U, Schmidt H, Karch H, Cheasty T, and Caprioli A. 2001. Detection and characterization of Shiga toxin-producing *Escherichia coli* in feral pigeons. *Vet Microbiol* 82:275-283.
191. Morgan AK, Piispanen, Humphreys, and Murphy D. 2005. A cluster of cases of haemolytic uraemic syndrome in north Queensland associated with a novel Shiga-like toxin-producing *Escherichia coli*. *Commun Dis Intelligence* 29:191-193.
192. Muraoka R, Okazaki M, Fujimoto Y, Jo N, Yoshida R, Kiyoyama T, Oura Y, Hirakawa K, Jyukurogi M, Kawano K, Okada M, Shioyama Y, Iryoda K, Wakamatsu H, Kawabata N. 2007. An enterohemorrhagic *Escherichia coli* O103 outbreak at a nursery school in Miyazaki Prefecture, Japan. *Jap J Infect Dis* 60:410-411.
193. Murphy M, Buckley JF, Whyte P, O'Mahony M, Anderson W, Wall PG, and Fanning S. 2007. Surveillance of dairy production holdings supplying raw milk to the farmhouse cheese sector for *Escherichia coli* O157, O26 and O111. *Zoonoses Public Health* 54:358-365.
195. Nagy JO, Zhang Y, Yi W, Liu X, Motari E, Song JC, Lejeune JT, and Wang PG. 2008. Glycopolymers as a chromatic biosensor to detect shiga-like toxin producing *Escherichia coli* O157:H7. *Bioorgan Med Chem Lett* 18:700-703.
196. ND Dept. of Health. Foodborne outbreaks in North Dakota 1988-2007. 2007. http://www.ndhealth.gov/disease/GI/Docs/Foodborne_Outbreaks_in_ND.pdf.
197. Nielsen EM, Jensen C, and Baggesen DL. 2005. Evidence of transmission of verocytotoxin-producing O111 from a cattle stable to a child. *Clin Microbiol Infect* 11:767-770.
198. Nielsen EM, Scheutz F, and Torpdahl M. 2006. Continuous surveillance of shiga toxin-producing *Escherichia coli* infections by pulsed-field gel electrophoresis shows that most infections are sporadic. *Foodborne Path Dis* 3:81-87.
199. Nielsen EM, Skov MN, Madsen JJ, Lodal J, Jespersen JB, and Baggesen DL. 2004. Verocytotoxin-producing *Escherichia coli* in wild birds and rodents in close proximity to farms. *Appl Environ Microbiol* 70:6944-6947.
200. Notario R, Fain JC, Prado V, Rios M, Borda N, and Gambande T. 2000. Animal reservoir and genotypic characterization of enterohemorrhagic *Escherichia coli* (EHEC) in Argentina. *Rev Med Chile* 128:1335-1341.

201. Novotna R, Alexa P, Hamrik J, Madanat A, Smola J, and Cizek A. 2005. Isolation and characterization Shiga toxin-producing *Escherichia coli* from sheep and goats in Jordan with evidence of multiresistant serotype O157:H7. *Vet Medicina* 50:111-118.
202. O'Hanlon KA, Catarama TMG, Blair IS, McDowell DA, and Duffy G. 2005. Comparison of a real-time PCR and an IMS/culture method to detect *Escherichia coli* O26 and O111 in minced beef in the Republic of Ireland. *Food Microbiol* 22:553-560.
203. Ogura Y, Ooka T, Iguchi A, Toh H, Asadulghani M, Oshima K, Kodama T, Abe H, Nakayama K, Kurokawa K, Tobe T, Hattori M, and Hayashi T. 2009. Comparative genomics reveal the mechanism of the parallel evolution of O157 and non-O157 enterohemorrhagic *Escherichia coli*. *Proc Nat Acad Sci USA* 106:17939-17944.
204. Oklahoma State Department of Health. Epidemiological investigation of restaurant-associated *Escherichia coli* O111:NM outbreak - Mayes County, Oklahoma; final report. 2009. 2009.
205. Oporto B, Esteban JI, Aduriz G, Juste RA, and Hurtado A. 2008. *Escherichia coli* O157:H7 and non-O157 shiga toxin-producing *E. coli* in healthy cattle, sheep and swine herds in northern Spain. *Zoonoses Public Health* 55:73-81.
206. Orden JA, Cortés C, Horcajo P, De La Fuente R, Blanco JE, Mora A, López C, Blanco J, Contreras A, Sánchez A, Corrales JC, and Domínguez-Bernal G. 2008. A longitudinal study of verotoxin-producing *Escherichia coli* in two dairy goat herds. *Vet Microbiol* 132:428-434.
207. Orth D, Grif K, Dierich MP, and Wurzner R. 2007. Variability in tellurite resistance and the *ter* gene cluster among shiga Toxin-producing *Escherichia coli* isolated from humans, animals and food. *Res Microbiol* 158:105-111.
208. Orth D, Grif K, Dierich MP, and Würzner R. 2006. Sorbitol-fermenting shiga toxin-producing *Escherichia coli* O157: indications for an animal reservoir. *Epidemiol Infect* 134:719-723.
209. OzFoodNet Working Group. 2003. OzFoodNet: Enhancing foodborne disease surveillance across Australia: quarterly report, July to September, 2002. *Commun Dis Intell* 27:90.
210. OzFoodNet Working Group. 2007. Monitoring the incidence and causes of diseases potentially transmitted by food in Australia: Annual report of the OzFoodNet Network, 2006. *Commun Dis Intell* 31:345-365.
211. OzFoodNet Working Group. 2009. OzFoodNet quarterly report, 1 October to 31 December 2008. *Commun Dis Intell* 33:53-58.
212. Palumbo SA, Pickard A, and Call LE. 1997. Population changes and verotoxin production of enterohemorrhagic *Escherichia coli* strains inoculated in milk and ground beef held at low temperatures. *J Food Prot* 60:746-750.
213. Park CH, Kim HJ, Hixon DL, and Bubert A. 2003. Evaluation of the Duopath Verotoxin Test for detection of shiga toxins in cultures of human stools. *J Clin Microbiol* 41:2650-2653.
214. Paton AW, Ratcliff RM, Doyle RM, Seymour-Murray J, Davos D, Lanser JA, and Paton JC. 1996. Molecular microbiological investigation of an outbreak of hemolytic-uremic syndrome caused by dry fermented sausage contaminated with shiga-like toxin-producing *Escherichia coli*. *J Clin Microbiol* 34:1622-1627.
215. Pedersen K, Clark L, Andelt WF, and Salman MD. 2006. Prevalence of shiga toxin-producing *Escherichia coli* and *Salmonella enterica* in rock pigeons captured in Fort Collins, Colorado. *J Wildlife Dis* 42:46-55.
216. Perelle S, Dilasser F, Grout J, and Fach P. 2005. Detection of *Escherichia coli* serogroup O103 by real-time

- polymerase chain reaction. *J Appl Microbiol* 98:1162-1168.
217. Perelle S, Dilasser F, Grout J, and Fach P. 2007. Screening food raw materials for the presence of the world's most frequent clinical cases of shiga toxin-encoding *Escherichia coli* O26, O103, O111, O145 and O157. *Int J Food Microbiol* 113:284-288.
 218. Perelle S, Dilasser F, Grout JL, and Fach P. 2004. Detection by 5'-nuclease PCR of shiga-toxin producing *Escherichia coli* O26, O55, O91, O103, O111, O113, O145 and O157:H7, associated with the world's most frequent clinical cases. *Molec Cell Probes* 18:185-192.
 219. Pichner R, Sander A, Steinruck H, and Gareis M. 2005. Occurrence of *Salmonella* spp. and shigatoxin-producing *Escherichia coli* (STEC) in horse faeces and horse meat products. *Berlin Munch Tierarzt Wochenschr* 118:321-325.
 220. Possé B, De Zutter L, Heyndrickx M, and Herman. 2008. Quantitative isolation efficiency of O26, O103, O111, O145 and O157 STEC serotypes from artificially contaminated food and cattle faeces samples using a new isolation protocol. *J Appl Microbiol* 105:227-235.
 221. Pradel N, Bertin Y, Martin C, and Livrelli V. 2008. Molecular analysis of shiga toxin-producing *Escherichia coli* strains isolated from hemolytic-uremic syndrome patients and dairy samples in France. *Appl Environ Microbiol* 74:2118-2128.
 222. Prager R, A. Fruth A, U. Siewert U, U. Strutz U, and H. Tschape H. 2009. *Escherichia coli* encoding shiga toxin 2f as an emerging human pathogen. *Int J Med Microbiol* 299:343-353.
 223. Rabatsky-Ehr T, Dingman D, Marcus R, Howard R, Kinney A, and Mshar P. 2002. Deer meat as the source for a sporadic case of *Escherichia coli* O157:H7 infection, Connecticut. *Emerg Infect Dis* 8:525-527.
 224. Ram S, Vajpayee P, Singh RL, and Shanker R. 2009. Surface water of a perennial river exhibits multi-antimicrobial resistant shiga toxin and enterotoxin producing *Escherichia coli*. *Ecotoxicol Environ Safety* 72:490-495.
 225. Reinders RD, Weber MF, Lipman LJA, Verhoeff J, and Bijker PGH. 2001. Control of VTEC in Dutch livestock and meat production. *Int J Food Microbiol* 66:79-83.
 226. Rey J, Blanco JE, Blanco M, Mora A, Dahbi G, Alonso JM, Hermoso M, Hermoso J, Alonso MP, Usera MA, Gonzalez EA, Bernardez MI, and Blanco J. 2003. Serotypes, phage types and virulence genes of shiga-producing *Escherichia coli* isolated from sheep in Spain. *Vet Microbiol* 94:47-56.
 227. Rey J, Sanche S, Blanco JE, De Mendoza JH, De Mendoza MH, Garcia A., Gil C., Tejero N, Rubio R, and Alonso JM. 2006. Prevalence, serotypes and virulence genes of shiga toxin-producing *Escherichia coli* isolated from ovine and caprine milk and other dairy products in Spain. *Int J Food Microbiol* 107:212-217.
 228. Rhoades JR, Duffy G, and Koutsoumanis K. 2009. Prevalence and concentration of verocytotoxigenic *Escherichia coli*, *Salmonella Enterica* and *Listeria monocytogenes* in the beef production chain: a review. *Food Microbiol* 26:357-376.
 229. Riley LW, Remis RS, Helgerson SD, McGee HB, Wells JG, Davis BR, Hebert RJ, Olcott ES, Johnson LM, Hargrett NT, Blake PA, and Cohen ML. 1983. Hemorrhagic colitis associated with a rare *Escherichia coli* serotype. *N Engl J Med* 308:681-685.
 230. Rivas M, Sosa-Estani S, Rangel J, Caletti MG, Valles P, Roldan CD, Balbi L, De Mollar MCM, Amoedo D, Miliwebsky E, Chinen I, Hoekstra RM, Mead P, and Griffin PM. 2008. Risk Factors for sporadic shiga toxin-producing *Escherichia coli* infections in children, Argentina. *Emerg Infect Dis* 14:763-771.

231. Sanchez S, Garcia-Sanchez A, Martinez R, Blanco J, Blanco JE, Blanco M, Dahbi G, Mora A, Hermoso De Mendoza J, Alonso JM, and Rey J. 2009. Detection and characterisation of shiga toxin-producing *Escherichia coli* other than *Escherichia coli* O157:H7 in wild ruminants. *Vet J* 180:384-388.
232. Scaife HR, Cowan D, Finney J, Kinghorn-Perry SF, and Crook B . 2006. Wild rabbits (*Oryctolagus cuniculus*) as potential carriers of verocytotoxin-producing *Escherichia coli*. *Vet Rec* 159:175-178.
233. Schamberger GP and Diez-Gonzalez F. 2004. Characterization of colicinogenic *Escherichia coli* strains inhibitory to enterohemorrhagic *Escherichia coli*. *J Food Prot* 67:486-492.
234. Schimmer B, Nygard K, Eriksen HM, Lassen J, Lindstedt BA, Brandal LT, Kapperud G, and Aavitsland P. 2008. Outbreak of haemolytic uraemic syndrome in Norway caused by Stx(2)-positive *Escherichia coli* O103:H25 traced to cured mutton sausages. *BMC Infectious Dis* 8.
235. Schmidt H, Scheef J, Morabito S, Caprioli A, Wieler LH, and Karch A. 2000. A new shiga toxin 2 variant (Stx2f) from *Escherichia coli* isolated from pigeons. *Appl Environ Microbiol* 66:1205-1208.
236. Schouten JM, van de Giessen AW, Frankena K, De Jong MCM, and Graat EAM. 2005. *Escherichia coli* O157 prevalence in Dutch poultry, pig finishing and veal herds and risk factors in Dutch veal herds. *Prev Vet Med* 70:1-15.
237. Sekse C, O'Sullivan K, Granum PE, Rørvik LM, Wasteson Y, and Jørgensen HJ. 2009. An outbreak of *Escherichia coli* O103:H25 - bacteriological investigations and genotyping of isolates from food. *Int J Food Microbiol* 133:259-264.
238. Sepehriseresht S, Salehi TZ, Sattari M, Tadjbakhsh H, and Aslani MM. 2009. Detection of shigatoxigenic *Escherichia coli* from fecal samples of calves and cattle by molecular and serological methods. *Comp Clin Pathol* 18:53-57.
239. Shaw DJ, Jenkins C, Pearce MC, Cheasty T, Gunn GJ, Dougan G, Smith HR, Woolhouse A, and Frankel G. 2004. Shedding patterns of verocytotoxin-producing *Escherichia coli* strains in a cohort of calves and their dams on a Scottish beef farm. *Appl Environ Microbiol* 70:7456-7465.
240. Shaw RK , Berger CN, Feys B, Knutton S, Pallen MJ, and Frankel G. 2008. Enterohemorrhagic *Escherichia coli* Exploits EspA filaments for attachment to salad leaves. *Appl Environ Microbiol* 74:2908-2914.
241. Sheng HQ, Knecht HJ, Kudva IT, and Hovde CJ. 2006. Application of bacteriophages to control intestinal *Escherichia coli* O157:H7 levels in ruminants. *Appl Environ Microbiol* 72:5359-5366.
242. Shere JA, Kaspar CW, Bartlett KJ, Linden SE, Norell B, Francey S, and Schaefer DM. 2002. Shedding of *Escherichia coli* O157:H7 in dairy cattle housed in a confined environment following waterborne inoculation. *Appl Environ Microbiol* 68:1947-1954.
243. Skaloud J, Pokludova L, Novotna R, and Cizek A. 2003. Evaluation by conductance assay of shiga toxin producing *Escherichia coli* (STEC) O157 and O26 and their sensitivity to selected disinfectants. *Acta Veterinaria Brno* 72:101-109.
244. Smith CJ, Olszewski AM, and Mauro SA. 2009. Correlation of shiga toxin gene frequency with commonly used microbial indicators of recreational water quality. *Appl Environ Microbiol* 75:316-321.
245. Smith DL, Wareing BM, Fogg PCM, Riley LM , Spencer M, Cox MJ, Saunders JR, McCarthy AJ, and Allison HE. 2007. Multilocus characterization scheme for shiga toxin-encoding bacteriophages. *Appl Environ Microbiol* 73:8032-8040.
246. Smith KE, Stenzel SA, Bender JB, Wagstrom E, Soderlund D, Leano FT, Taylor CM, Belle-Isle PA, and

- Danila R. 2004. Outbreaks of enteric infections caused by multiple pathogens associated with calves at a farm day camp. *Pediatr Infect Dis J* 23:1098-1104.
247. Sonntag AK, Zenner E, Karch H, and Bielaszewska M. 2005. Pigeons as a possible reservoir of shiga toxin 2f-producing *Escherichia coli* pathogenic to humans. *Berlin Munch Tierarz Wochenschr* 118:464-470.
248. Sonoda C, Tagami A, Nagatomo D, Yamada S, Fuchiwaki R, Haruyama M, Nakamura Y, Kawano K, Okada M, Shioyama Y, Iryoda K, Wakamatu H, and Hidaka Y. 2008. An enterohemorrhagic *Escherichia coli* O26 outbreak at a nursery school in Miyazaki, Japan. *Jap J Infect Dis* 61:92-93.
249. Stephan R, Schumacher S, Corti S, Krause G, Danuser J, and Beutin L. 2008. Prevalence and characteristics of shiga toxin-producing *Escherichia coli* in Swiss raw milk cheeses collected at producer level. *J Dairy Sci* 91:2561-2565.
250. Tarawneh KA, Al-Tawarah NM, Abdel-Ghani AH, Al-Majali AM, and Khleifat KM. 2009. Characterization of verotoxigenic *Escherichia coli* (VTEC) isolates from faeces of small ruminants and environmental samples in southern Jordan. *Jf Basic Microbiol* 49:310-317.
251. Tarr CL, Nelson AM, Beutin L, Olsen KEP, and Whittam TS. 2008. Molecular characterization reveals similar virulence gene content in unrelated clonal groups of *Escherichia coli* of serogroup O174 (OX3). *J Bacteriol* 190:1344-1349.
252. Teel LD, Daly JA, Jerris RC, Maul D, Svanas G, O'Brien AD, and Park CH. 2007. Rapid detection of shiga toxin-producing *Escherichia coli* by optical immunoassay. *J Clin Microbiol* 45:3377-3380.
253. Thompson LH, Giercke S, Beaudoin C, Woodward D, and Wylie JL. 2005. Enhanced surveillance of non-O157 verotoxin-producing *Escherichia coli* in human stool samples from Manitoba. *Can J Infect Dis Med Microbiol* 16:329-334.
254. Tkalcic S, Zhao T, Harmon BG, Doyle MP, Brown CA, and Zhao P. 2003. Fecal shedding of enterohemorrhagic *Escherichia coli* in weaned calves following treatment with probiotic *Escherichia coli*. *J Food Prot* 66:1184-1189.
255. Urdahl AM, Beutin L, Skjerve E, and Wasteson Y. 2002. Serotypes and virulence factors of shiga toxin-producing *Escherichia coli* Isolated from healthy Norwegian sheep. *J Appl Microbiol* 93:1026-1033.
256. Valcour JE, Michel P, McEwen SA, and Wilson JB. 2002. Associations between indicators of livestock farming intensity and incidence of human shiga toxin-producing *Escherichia coli* infection. *Emerg Infect Dis* 8:252-257.
257. Van Duynhoven YTHP, Friesema IHM, Schuurman T, Roovers A, Van Zwet AA, Sabbe LJM, Van Der Zwaluw WK, Notermans DW, Mulder B, Van Hannen EJ, Heilmann FGC, Buiting A, Jansen R, and Kooistra-Smid AMD. 2008. Prevalence, characterisation and clinical profiles of shiga toxin-producing *Escherichia coli* in the Netherlands. *Clin Microbiol Infect* 14:437-445.
258. Vanselow BA, Krause DO, and McSweeney CS. 2005. The shiga toxin-producing *Escherichia coli*, their ruminant hosts, and potential on-farm interventions: a review. *Austral J Agr Res* 56:219-244.
259. Venter P, Abraham M, Lues JFR, and Ivanov I. 2006. The influence of sanitizers on the lipopolysaccharide composition of *Escherichia coli* O111. *Int J Food Microbiol* 111:221-227.
260. Vettorato MP, De Castro AFP, Cergole-Novella MC, Camargo FLL, Irino K, and Guth BEC. 2009. Shiga toxin-producing *Escherichia coli* and atypical enteropathogenic *Escherichia coli* strains isolated from healthy sheep of different populations in Sao Paulo, Brazil. *Lett Appl Microbiol* 49:53-59.

261. Vimont A, Delignette-Muller ML, and Vernozy-Rozand C. 2007. Supplementation of enrichment broths by novobiocin for detecting shiga toxin-producing *Escherichia coli* from food: a controversial use. *Lett Appl Microbiol* 44:326-331.
262. Vimont A, Vernozy-Rozand C, and Delignette-Muller ML. 2006. Isolation of *E. coli* O157:H7 and non-O157 STEC in different matrices: review of the most commonly used enrichment protocols. *Lett Appl Microbiol* 42:102-108.
263. Viscardi M, Perugini AG, Auriemma C, Capuano F, Morabito S, Kim KP, Loessner MJ, and G. Iovane G. 2008. Isolation and characterisation of two novel coliphages with high potential to control antibiotic-resistant pathogenic *Escherichia coli* (EHEC and EPEC). *Int J Antimicrob Agents* 31:152-157.
264. Vojdani JD, Beuchat LR, and Tauxe RV. 2008. Juice-associated outbreaks of human illness in the United States, 1995 through 2005. *J Food Prot* 71:356-364.
265. Von Muffling T, Smajilovic M, Nowak B, Sammet K, Bulte M, and Klein G. 2007. Preliminary study of certain serotypes, genetic and antimicrobial resistance profiles of verotoxigenic *Escherichia coli* (VTEC) isolated in Bosnia and Germany from cattle or pigs and their products. *Int J Food Microbiol* 117:185-191.
266. Watterworth L, Rosa B, Schraft H, Topp E, and Leung KT. 2006. Survival of various ERIC-genotypes of shiga toxin-producing *Escherichia coli* in well water. *Water Air Soil Pollut* 177:367-382.
267. Wells JG, Davis B R, Wachsmuth IK, Riley LW, Remis RS, Sokolow R, and Morris GK. 1983. Laboratory investigation of hemorrhagic colitis outbreaks associated with a rare *Escherichia coli* serotype. *J Clin Microbiol* 18:512-20.
268. Werber D, Beutin L, Pichner R, Stark K, and Fruth A. 2008. Shiga toxin-producing *Escherichia coli* serogroups in food and patients, Germany. *Emerg Infect Dis* 14:1803-1806.
269. Werber D, Frank C, Wadl M, Karch H, Fruth A, and Stark K. 2008. Looking for tips to find icebergs--surveillance of haemolytic uraemic syndrome to detect outbreaks of shiga toxin-producing *E. coli* infection. *Euro Surveill* 13:pii=8053.
270. Werber D, Fruth A, Liesegang A, Littmann M, Buchholz U, Prager R, Karch H, Breuer T, Tschape H, and Ammon A. 2002. A multistate outbreak of Shiga toxin-producing *Escherichia coli* O26:H11 infections in Germany, detected by molecular subtyping surveillance. *J Infect Dis* 186:419-422.
271. Widiasih DA, Ido N, Omoe K, Sugii S, and Shinagawa K. 2004. Duration and magnitude of faecal shedding of Shiga toxin-producing *Escherichia coli* from naturally infected cattle. *Epidemiol Infect* 132:67-75.
272. Willford J, Mills K, and Goodridge LD. 2009. Evaluation of three commercially available enzyme-linked immunosorbent assay kits for detection of shiga toxin. *J Food Prot* 72:741-747.
273. Zaki MES and El-Adrosy H. 2007. Diagnosis of shiga toxin producing *Escherichia coli* infection, contribution of genetic amplification technique. *Microbes Infect* 9 :200-203.
274. Zhang WL, Bielaszewska M, Pulz M, Becker K, Friedrich AW, Karch H, and Kuczius T. 2008. New immuno-PCR assay for detection of low concentrations of shiga toxin 2 and its variants. *J Clin Microbiol* 46:1292-1297.
275. Zhao T, Zhao P, West JW, Bernard JK, Cross HG, and Doyle MP. 2006. Inactivation of enterohemorrhagic *Escherichia coli* in rumen content- or feces-contaminated drinking water for cattle. *Appl Environ Microbiol* 72:3268-3273.
276. Zweifel C, Blanco JE./Blanco M, Blanco J, and Stephan R. 2004. Serotypes and virulence genes of ovine

- non-0157 shiga toxin-producing *Escherichia coli* in Switzerland. Int J Food Microbiol 95:19-27.
277. Zweifel C, S. Schumacher, Beutin L, Blanco J, and Stephan R . 2006. Virulence profiles of Shiga toxin 2e-producing *Escherichia coli* isolated from healthy pig at slaughter. Vet Microbiol 117:328-332.
278. Zweifel C, Zychowska MA, and Stephan R. 2004. Prevalence and characteristics of shiga toxin-producing *Escherichia coli*, *Salmonella* spp. and *Campylobacter* spp. isolated from slaughtered sheep in Switzerland. Int J Food Microbiol 92:45-53.